


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/64		A2	(11) International Publication Number: WO 96/31612
			(43) International Publication Date: 10 October 1996 (10.10.96)
(21) International Application Number: PCT/US96/04688		(74) Agent: MUELLER, Lisa, V.; Greer, Burns & Crain, Ltd., Sears Tower, Suite 8660, 233 South Wacker Drive, Chicago, IL 60606 (US).	
(22) International Filing Date: 5 April 1996 (05.04.96)			
(30) Priority Data: 08/417,618 6 April 1995 (06.04.95) US		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(60) Parent Application or Grant (63) Related by Continuation US 08/417,618 (CIP) Filed on 6 April 1995 (06.04.95)			
(71) Applicant (for all designated States except US): SEMINIS VEGETABLES [US/US]; P.O. Box 4206, Saticoy, CA 93007-4206 (US).		Published Without international search report and to be republished upon receipt of that report.	
(72) Inventors; and (75) Inventors/Applicants (for US only): TRULSON, Anna, Julia [US/US]; 2751 Concord Avenue, Davis, CA 95616 (US). GREEN, Charles, Edward [US/US]; 39648 Lupine Court, Davis, CA 95616 (US). BRAUN, Carl, Joseph, III [US/US]; 616 Spring Court, Woodland, CA 95776 (US).			

(54) Title: PROCESS FOR SELECTION OF TRANSGENIC PLANT CELLS

(57) Abstract

The present invention involves a selection process for increasing the number of transformed plant cells from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions. In accordance with this process, a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and growth by non-transformed plant cells except for a source of carbon that supports growth and proliferation and about 1.5 to 3 times the standard amount of phosphorus. The source of carbon utilized is replaced by an encrypted carbon source that does not support growth and proliferation of said non-transformed cells. The transformed plant cells have a heterologous DNA segment inserted into their genome that contains at least one expression cassette. The expression cassette contains a heterologous DNA selectable marker segment that includes a first heterologous gene that encodes a heterologous enzyme that on expression converts the encrypted carbon source into a carbon source that supports growth and proliferation by the transformed plant cells under heterotrophic culture conditions. Additionally, the present invention also contemplates a selection process for increasing the number of transformed plant cells from a mixture of transformed and non-transformed plant cells cultured under delayed selective culture conditions.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CZ	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

PROCESS FOR SELECTION
OF TRANSGENIC PLANT CELLS

5

Description

Technical Field

The present invention relates to genetically transformed plants, and more particularly to a process for the selection of transformed plant cells cultured under heterotrophic conditions, a process for selectively increasing the number of transformed plants regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions, as well as to transformed rooted plants and a kit for forming transformed plants.

Background Art

Techniques for selection of transgenic plant cells that are used currently involve the utilization of prokaryotic genes conferring resistance to toxic substances that are added to the regeneration medium, but that themselves are not essential for cell growth. These toxic additives are either antibiotics, herbicides, amino acids or amino acid analogs added in toxic concentrations. Although generally effective in selecting genetically modified plant cells, these systems have limitations. For example, they may not be effective in some types of plants, or they may result in abnormal phenotypes of the genetically engineered plants. Additionally, the use of these selective agents can have undesirable side effects and thus have raised public concern about their environmental safety.

Antibiotics such as kanamycin, G418, hygromycin, bleomycin and streptomycin, among others, have been used for selection of transgenic plants [Bevan et al., Nature, 394:184-187 (1983); Dekeyser et al., Plant Phys., 90:217-223 (1989); Hille et al.,

Plant Mol. Biol., 7:171-176 (1986); Jones et al., Mol. Gen. Gen., 210:86-91 (1987); Mulsant et al., Som. Cell Mol. Gen., 14:243-252 (1988); Van den Elzen et al., Plant Mol. Biol., 5:299-302 (1985); Waldron et al., Plant Mol. Biol., 5:103-108 (1985)]. However, not all plants are equally sensitive to certain antibiotics. Kanamycin, which is used commonly, is not effective as a selective agent in gramineaceous plants; some plants from this group can tolerate up to 800 mg/L of kanamycin [Dekeyser et al., Plant Phys., 90:217-223 (1989); Hauptmann et al., Plant Physiol., 86:1216-1222 (1988)].

Additionally, there are negative aspects of antibiotic use, primarily, but not exclusively, due to their high toxicity against eukaryotic cells. For example, hygromycin has a broad spectrum of toxicity against prokaryotic and eukaryotic cells due to interference with protein synthesis [Waldron et al., Plant Mol. Biol., 5:103-108 (1985)]. Bleomycin is a cytotoxic drug used in human cancer therapy and has known pulmonary toxicity [Gatignol et al., FEBS Lett., 230:171-175 (1988); Hille et al., Plant Mol. Biol., 7:171-176 (1986); Mulsant et al., Som. Cell Mol. Gen., 14:243-252 (1988)]. Many antibiotics are also potent allergens. Thus, extensive use of antibiotics in routine selections of transgenic plants can pose a health hazard to the research staff. It is also possible that upon lysis of transgenic plant cells, the DNA encoding for the antibiotic resistance may be taken up by bacteria, rendering them antibiotic-resistant and thereby a threat to the public health.

Herbicides such as chlorsulfuron, 2,4-D, glyphosate, phosphinotricin, and others, have been proposed as selective agents [De Block et al., EMBO J., 6:2513-2518 (1987); Dekeyser et al., Plant Phys., 90:217-223 (1989); Li et al., Plant Phys., 100:662-668 (1992); Streber et al., Bio/Technology, 7:811-816 (1989); Shah et al., Science, 233:478-481 (1986); White

et al., Nucl. Acid Res., 18:1062 (1990)]. The use of herbicide resistance in identification of transgenic plants can result in increased weediness of transgenic plants because they can become herbicide-resistant weeds in the alternate years of crop rotation. Out-crossing to wild relatives is also a concern. If the mechanism of resistance to the herbicide is based on detoxification, there is a potential for one or more of its metabolites to be toxic (Stalker, et al., Science, 242:419-423 (1988)). Further, the use of herbicide-resistant crops probably will increase the herbicide load in the environment.

Certain amino acids such as lysine and threonine, or the lysine derivative amino ethyl cysteine (AEC), can also be used as selective agents due to their ability to inhibit cell growth when applied at high concentrations [Shaul et al., The Plant J., 2:203-209 (1992); Perl et al., Bio/Technology, 11:715-718 (1993)]. In this selection system, expression of the selectable marker gene, which permits the transgenic cells to grow under selection, results in overproduction of specific amino acids by transgenic cells, which counteracts the selective pressure. In some cases, this results in abnormal plant development [Shaul et al., The Plant J., 2:203-209 (1992)]. Moreover, transgenic plants that are selected in such a system have permanently altered amino acid composition, which may affect the nutritional value of the transgenic plants [Perl et al., Bio/Technology, 11:715-718 (1993)].

To be generally useful, a selectable marker must meet certain criteria. Selection must be stringent with a minimum of non-transformed plant tissue escaping the selection process. The selection should result in a large number of independent transformation events and not significantly interfere with regeneration. In addition, the marker should work well in a large number of plant species, and there

should be an assay to score transformed tissue for confirmation that the marker gene is being expressed.

Another phenomenon associated with selectable markers is that once transformed plant cells are selected by means of the marker and rooted plants are regenerated, the marker gene continues to be expressed in the mature, autotrophically-grown plants. Thus, as was noted before, a transgenic plant can permanently express an altered amino acid composition or a gene for antibiotic resistance. It would be beneficial if once selected and regenerated, the autotrophic rooted plants expressed the marker to a lessened extent or not at all.

The present invention provides a process for selecting transformed plant cells that meets the above criteria, while avoiding toxic or environmentally unsafe substances. The present invention also provides a process for selectively increasing the number of transformed plant cells regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions. The present invention further provides a means by which expression of the selectable marker can be lessened or shut off after the selection process.

25

Brief Summary of the Invention

The present invention contemplates a process for selectively growing transformed plant cells cultured under heterotrophic conditions. Also contemplated is a process for selectively increasing the number of transformed plant cells regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions. Still further contemplated is a method for increasing the number of transformed plant cells regenerated from a mixture of transformed and non-transformed plant cells under delayed heterotrophic culture conditions. Still even further contemplated are transformed plants

whose genome contains an identifiable heterologous, exogenously supplied DNA segment that contains at least one expression cassette. A kit useful for transforming plant cells is also contemplated.

5 Thus, in one embodiment, a selection process for transformed plant cells is contemplated. In accordance with this process,

 (a) a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for growth and proliferation by those plant cells except for a source of carbon that is utilized to support that growth and proliferation. The source of carbon utilized is replaced by an encrypted or latent (growth-limiting) carbon source that does not support growth and proliferation by the non-transformed cells. The transformed cells of the mixture contain a genomic heterologous DNA segment that contains at least two expression cassettes,

10

15

20 The first expression cassette contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts the encrypted carbon source into a carbon source that supports growth and proliferation by the transformed plant cells under heterotrophic culture conditions. The first gene is operatively linked to (ii) a first promoter DNA segment that controls expression of the heterologous gene, and (iii) a termination DNA segment.

25

30 The second expression cassette contains (i) a second gene that is expressed in a transformed plant, and that gene is operatively linked to (ii) a second promoter DNA segment that controls expression of that second gene and (iii) a termination DNA segment.

35 (b) The heterotrophic culture conditions are maintained for a time period sufficient for the transformed plant cells to express the heterologous enzyme, proliferate and grow. Inasmuch as the non-

-6-

transformed plant cells cannot utilize the encrypted or latent carbon source, those cells do not grow and proliferate. The transformed cells that do grow and proliferate can thereby be selected from the non-transformed cells.

A particularly preferred first gene encodes the enzyme phosphomannose isomerase (*pmi*; EC 5.3.1.8) that converts non-utilizable mannose-6-phosphate into fructose-6-phosphate that can be used by plant cells as a carbon source to support cell growth and proliferation. The *pmi* gene is also known as *manA*, and this gene is often referred to herein as *pmi/manA*. The encrypted (growth-limiting) carbon source useful with this first gene is mannose. Another preferred useful gene encodes mannitol-1-oxidoreductase that converts mannitol into mannose, and here, mannitol is the encrypted (growth-limiting) carbon source. This second gene and its encrypted carbon source are used in plant cells that have previously been transformed with a *pmi/manA* gene. Another preferred first gene encodes human L-iditol dehydrogenase (EC 1.1.1.14) that converts sorbitol into fructose, so that sorbitol is used as the encrypted (growth-limiting) carbon source. Similar aldehyde reductase enzyme genes can also be used.

The proliferating cells so produced and selected can thereafter be harvested or regenerated by culture in appropriate media into mature plants via meristematic tissue or embryos, or via callus tissue conversion into meristematic tissue or embryos. Thus, the selected proliferating cells are preferably collected and thereafter regenerated into mature plants that grow autotrophically. The above process therefore more preferably utilizes the added steps of:

(c) recovering the selected proliferating cells; and/or

(d) regenerating plants from those proliferating cells.

The promoter of the first expression cassette is repressed by a product of the normal autotrophic metabolism of the transgenic plant, which product is also present in a non-transgenic plant. Exemplary preferred promoters include the cucumber isocitrate lyase promoter and the rice α -amylase Amy3A promoter.

The second gene and the genes in the subsequent cassettes can be any gene desired to be expressed in a plant, and its promoter and termination DNA segments can be any desired promoter and terminator that operate in plants.

In a second embodiment, a selection process for increasing the number of transformed plant cells from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions is contemplated. In accordance with this process,

(a) a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and growth by non-transformed plant cells except for a source of carbon that supports growth and proliferation and about 1.5 to 3 times the standard amount of phosphorus. The source of carbon utilized is replaced by an encrypted carbon source that does not support growth and proliferation of said non-transformed cells. The transformed cells of the mixture have a heterologous DNA segment inserted into their genome that contains at least one expression cassette.

At least one expression cassette contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts the encrypted carbon source into a carbon source that supports growth and proliferation by the transformed plant cells under heterotrophic culture conditions. The first gene is operatively linked to (ii) a first promoter DNA segment that controls

expression of the heterologous gene, and (iii) a termination DNA segment.

5 (b) The heterotrophic culture conditions are maintained for a time period sufficient for the transformed plant cells to express the heterologous enzyme, proliferate and grow.

10 In a third embodiment, a selection process for increasing the number of transformed plant cells from a mixture of transformed and non-transformed plant cells cultured under delayed selective culture conditions is contemplated. In accordance with this process,

15 (a) a mixture of transformed and non-transformed plant cells is cultured for up to two weeks in a first culture medium that contains the minimal nutrients required for proliferation and growth by both, transformed and non-transformed plant cells including a source of carbon that supports growth and proliferation of both the transformed and non-transformed plant cells. The transformed plant cells
20 have a heterologous genomic DNA segment that contains at least one expression cassette.

At least one expression cassette contains a heterologous DNA selectable marker segment that
25 includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts an encrypted carbon source into a carbon source that supports growth and proliferation of said transformed plant cells under heterotrophic culture conditions, said first gene being operatively linked to (ii) a
30 first promoter DNA segment that controls expression of said heterologous gene, and (iii) a termination DNA segment.

35 (b) After sufficient time in the first medium, the transformed and non-transformed plant cells are removed from the first culture medium.

(c) The transformed and non-transformed plant cells are then placed under heterotrophic culture

conditions in a second culture medium that contains the minimal nutrients required for proliferation and growth of the non-transformed plant cells except for an encrypted carbon source that does not support growth and proliferation of said non-transformed plant cells and 1.5 to 3 times the standard amount of phosphorous.

(d) The heterotrophic culture conditions are maintained for a time period sufficient for said transformed plant cells to express said heterologous enzyme, proliferate and grow.

A transgenic plant whose genome comprises a heterologous DNA segment that contains at least two expression cassettes is further contemplated.

The first expression cassette contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression during heterotrophic culture of cells from the transformed plant converts an encrypted carbon source that does not support growth and proliferation of non-transformed plant cells of the same type into a carbon source that supports growth and proliferation of those transformed cells. That first gene is operatively linked to (ii) a promoter DNA segment that controls expression of the heterologous gene and (iii) a termination DNA segment. The second expression cassette contains (i) a second gene that is expressed in the transformed plant that is operatively linked to (ii) a second promoter DNA segment that controls expression of the second gene and (iii) a termination DNA segment.

The before-noted preferences are also followed for the first gene and its promoter in the transgenic plant. The second gene and its promoter are also as discussed before.

A kit for forming transformed plant cells is also contemplated. That kit comprises:

(a) a first package containing a DNA segment for transforming plant cells that contains an

expression cassette operatively linked to a linker segment containing at least one restriction endonuclease site. The expression cassette contains a heterologous DNA selectable marker segment that

5 includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression during heterotrophic culture of transformed plant cells converts an encrypted carbon source that does not support growth and proliferation of non-transformed

10 plant cells into a carbon source that supports growth and proliferation the transformed cells. The first heterologous gene is operatively linked to (ii) a promoter DNA segment that controls expression of the heterologous gene and (iii) a termination DNA segment.

15 (b) a second package is also present that contains minimal nutrients required for growth and proliferation of plant cells during heterotrophic culture except for a source of carbon and about 1.5 to 3 times the standard amount of phosphorous. That

20 source of carbon is replaced by an encrypted carbon source that does not support growth and proliferation of non-transformed plant cells but does support growth and proliferation of a transformed plant cell whose genome contains the DNA segment of the first package.

25 Instructions for use of the kit components are also preferably provided.

The present invention has several benefits and advantages.

30 One benefit of the invention is that its selective growth process does not rely upon potentially harmful antibiotics, herbicides or other possibly toxic materials.

One advantage of the invention is that its encrypted (growth-limiting) carbon source can be and

35 preferably is a carbohydrate.

Another benefit of the invention is that expression of the selectable marker can be repressed in

-11-

the regenerated plant under autotrophic growth conditions.

Another advantage of the invention is that the selectable marker gene can be used with any second expressed gene.

5 Still another benefit of the invention is that the kit provides a ready means for inserting a second expression cassette into a plant transforming vector and an appropriate selection medium for the enhanced transformation and selected growth of
10 transformed plant cells.

Still another advantage is that successive transformations can be made in which one encrypted (growth-limiting) carbohydrate can be converted by a second selectable marker gene into another encrypted
15 (growth-limiting) carbohydrate that is converted into a useful carbon source by a first selectable marker gene also present in the transformed cells.

Still further benefits and advantages of the present invention will be apparent to the skilled
20 worker from the disclosure that follows.

Brief Description of the Sequence Listing

SEQ ID NO:1 is a sequence listing for the 1650 base pair Clal fragment that contains the
25 *Salmonella typhimurium* phosphomannose isomerase (*pmi*) gene.

SEQ ID NO:2 is the amino acid sequence for *pmi*.

SEQ ID NO:3 is the sequence of one of the
30 primers used to facilitate the cloning of the *pmi* gene into a plant transformation vector. The primer corresponds to bases 391 to 410 of SEQ ID NO:1 with a substitution of an adenine for a thymine at base 403.

SEQ ID NO:4 is the sequence of the other
35 primer used to facilitate the cloning of the *pmi* gene into a plant transformation vector. The primer corresponds to a complementary sequence to bases 1617 to 1641 of SEQ ID NO:1, with the substitution of a

-12-

cytosine for a guanine at base 1628 and a cytosine for thymine at base 1630.

Brief Description of the Drawings

5 Fig. 1 shows the relationship between mannitol and sucrose biosynthesis in plants. Circled numbers identify enzymes that catalyze the reaction shown as follows: (1) mannitol-1-phosphate phosphatase, (2) mannose kinase, (3) mannose-6-phosphate reductase,
10 (4) mannose-6-phosphate isomerase, and (5) sucrose phosphate synthase.

 Fig. 2 illustrates the construction of plasmid vector pET0148. The recombinant DNAs manipulated and produced by the construction process
15 are indicated in the figure by the circles. The construction proceeds by a series of steps as indicated by the arrows connecting the circles in the figure and as described in detail herein. Landmark and utilized restriction enzyme recognition sites are indicated on
20 the circles by labeled lines intersecting the circles. The relative location of individual genetic elements and their orientation are indicated by the labeled arrows inside the circles.

 Fig. 3 shows the relative NPTII expression in
25 populations of plants selected on kanamycin and mannose with plants regenerated with no selection. NPTII expression was determined by an ELISA, and the units of expression are absorbance values at 405 nM. Data for non-transformed plants (negative control) are shown as
30 closed circles. Data for use of kanamycin (positive control) for selection of transformed plants are shown as closed triangles. Data for the transformed plants selected on mannose are shown as closed squares.

35

Detailed Description of the InventionA. Definition of Terms

Amino Acid: All amino acid residues identified herein are in the natural L-configuration.

5 Autologous: A DNA segment or protein normally present in a non-transformed cell.

Explant: A piece of plant tissue.

Expression: The combination of intracellular processes, including transcription and translation undergone by a protein-encoding gene to produce a polypeptide.

10 Expression cassette: A DNA segment construct comprising a gene to be expressed operatively linked to a promoter DNA and to a DNA termination segment and sequences sufficient for translation, as well as any other regulatory signals needed to effect proper processing of the expression product.

15 Expression vector: A DNA sequence that forms control elements that regulate expression of a desired gene when operatively linked to that gene within a vector. An expression vector of particular interest also contains DNA segments that permit integration into the plant genome.

20 Gene: A sequence of nucleotides in the genome of an organism to which a specific function can be ascribed.

Heterologous: A DNA segment or protein not present in a non-transformed cell.

Integrated: A heterologous DNA sequence incorporated into a host chromosome is integrated.

Operatively linked or inserted: A first DNA sequence such as a promoter DNA sequence is operatively linked to a second DNA sequence such as a heterologous gene DNA sequence if the two are situated so that the promoter DNA sequence influences the transcription or translation of the heterologous gene DNA sequence.

35 Promoter: A recognition site on a DNA sequence or group of DNA sequences that provide an

-14-

expression control element for a gene and to which RNA polymerase specifically binds and initiates RNA synthesis (transcription) of that gene.

5 Protein-encoding gene: A DNA sequence that through an RNA intermediate encodes for a polypeptide; i.e., an amino acid residue sequence.

Recombinant DNA molecule: A hybrid DNA sequence comprising at least two nucleotide sequences not that are created *in vitro*.

10 Same type or same strain plant: A plant of the same cross as or a clone of the untransformed plant.

Transgenic plant: A plant that contains chromosomally integrated heterologous or foreign DNA whose expression may be constitutive or regulatable.

15 Vector: A DNA molecule capable of replication in a cell and/or to which another DNA segment can be operatively linked so as to bring about replication of the attached segment. A plasmid is an exemplary
20 vector.

B. Introduction

Higher plants (plants) in their adult
25 (mature) forms are autotrophic organisms. Thus, mature plants use carbon dioxide for most of their carbon requirements. When grown from seed, the germinating seedling initially uses part of the nutrient rich seed to supply its carbon requirements until it is capable
30 of carbon fixation. Similarly, when plant cells are grown in a culture medium, as cell suspensions, microspores, protoplasts or as explants (collectively referred to as plant cells in that each is from a plant and has a nucleus and cytoplasm surrounded by a
35 membrane), the cultured plant cells are heterotrophic and require the provision of an external source of carbon as well as other nutrients. That carbon source is typically sucrose, glucose or another carbohydrate

-15-

that can be readily metabolized by the growing cells so that they can grow, proliferate and differentiate, and then be regenerated into mature plants.

5 However, not all sources of carbon, including some carbohydrates, can be metabolized by cultured plant cells. Cultured plants do not possess all of the necessary enzymes needed to convert many simple organic molecules into sources of carbon that can support growth and proliferation under tissue culture
10 conditions.

For example, mannose, a saccharide that is present in many mammalian glycoproteins cannot be utilized as a carbon source for many cultured plant cells. Thus, as is illustrated hereinafter, mannose is
15 taken up by cultured tomato cells, for example, and is converted into mannose-6-phosphate, but mannose-6-phosphate cannot be utilized by tomato plant cells as a source of carbon for cell growth and proliferation.

The present invention utilizes the fact that
20 plant cells cannot grow and proliferate using many small, carbon-containing compounds as a source of carbon during heterotrophic culture as a means of selectively growing transformed (genetically engineered) plant cells. This is accomplished, inter
25 alia, by using a marker gene for cell transformation that converts a source of carbon that does not support cell growth and proliferation (a non-useful carbon source) into a carbon source that supports cell growth and proliferation (a useful carbon source).

30 Thus, when a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions on a medium that contains a non-useful source of carbon that does not support cell growth and proliferation of non-transformed plant
35 cells, referred to herein as an encrypted, latent or growth-limiting carbon source, only those plant cells grow and proliferate that are transformed with a selectable marker gene whose expressed product converts

-16-

the encrypted carbon source into a useful source of carbon. This carbon source is encrypted (in code) because the plant cells cannot use it to grow and proliferate, and it is only after the selectable marker gene product that is present in the transformed callus decodes (decrypts) or acts upon that carbon source does it become useful to support growth and proliferation in the transformed plant cells.

10 C. The Processes of the Invention

In one embodiment, a selection process for transformed plant cells is contemplated. In accordance with this process,

15 (a) a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and differentiation by those plant cells except for a useful source of carbon. The normally present useful
20 source of carbon is replaced by an encrypted, latent or growth-limiting carbon source that does not support growth and proliferation of the non-transformed cells. The transformed cells of the mixture contain a genomic (chromosomally integrated) heterologous DNA segment
25 that contains two expression cassettes.

 The first expression cassette contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts the
30 encrypted carbon source into a useful carbon source that supports growth and proliferation of the transformed plant cells under heterotrophic culture conditions. That first heterologous gene is operatively linked to (ii) a first promoter DNA segment
35 that controls expression of the heterologous gene, and (iii) a termination DNA segment. The second expression cassette contains (i) a second heterologous gene that is expressed in a transformed plant, and that second

heterologous gene is operatively linked to (ii) a second promoter DNA segment that controls expression of the second heterologous gene and (iii) a termination DNA segment.

5 (b) The heterotrophic culture conditions are maintained for a time period sufficient for the transformed plant cells to express the heterologous enzyme, grow and proliferate. The non-transformed plant cells do not utilize the encrypted or latent
10 carbon source and do not grow and proliferate as well as the transformed cells. The transformed cells that do grow and proliferate are thereby selected away from the non-transformed cells, and are preferably collected for regeneration into plants.

15 The first expression cassette DNA thus contains three operatively linked elements: (1) a first heterologous gene that encodes a heterologous enzyme that converts an encrypted carbon source into a useful carbon source, (2) a first promoter DNA segment that
20 controls the expression of the heterologous gene, and (3) a first termination DNA segment that causes termination of transcription and polyadenylation of the 3'-end of the translated RNA.

In some cases, a gene that is functionally
25 similar to the first gene of the first expression cassette is present in the genome of the non-transformed plant. That endogenous gene is, however, functionally not present in the heterotrophically cultured plant cells inasmuch as the non-transformed
30 plant cells do not grow and proliferate using the same encrypted carbon source that the heterologous gene product of that first gene converts into a useful carbon source. This concept is exemplified by the fact that tomato plants do not grow and proliferate when
35 heterotrophically cultured using either lactose or salicin as carbon sources. Lactose is susceptible to cleavage by a β -galactosidase into glucose and galactose, whereas salicin can be cleaved by a β -

glucosidase to form glucose. Nonetheless, one commercially available β -galactosidase is recovered from jack beans, whereas a commercially available β -galactosidase is recovered from sweet almonds (Sigma Chemical Co.), indicating that the plant genomes contain genes for each enzyme that are not sufficiently expressed at all times.

The choice of the first gene depends upon its encoded enzyme and the function of that enzyme, which, in this invention, is to convert an encrypted, non-useful carbon source into a useful carbon source that supports the growth and proliferation of heterotrophically-cultured transgenic plant cells. Continuing this choice of the first gene backwards, one assays carbon sources that the plant cells to be transformed cannot utilize to grow and proliferate.

Such an assay is readily carried out by culturing the desired non-transformed plant cells in a medium that contains minimal nutrients required for cell growth and proliferation except for a source of carbon. Such media are well known in the art and are exemplified by the well known medium of Murashige and Skoog, Physiol. Plant, 15:437-498(1962); MS salts. The glucose or sucrose usually present in that medium is replaced with the carbon source to be assayed that is present at about the same concentration as was the glucose or sucrose.

The plant cells are then cultured in the resulting medium following usual procedures. An assayed carbon source that does not support growth and proliferation of the heterotrophically cultured cells is determined. Exemplary carbon sources that do not support the growth and proliferation of the illustrative tomato cells used herein include mannose, mannitol, sorbitol, lactose, trehalose and salicin. In addition, oat, maize, melon and squash cells are shown not to grow and proliferate using mannose as the sole carbon source.

An enzyme is then selected that converts the non-utilizable carbon source into a carbon source that can be utilized by the plants. Exemplary sources of carbon that can be used by plant cells to support growth and proliferation are the compounds of the Calvin cycle and include glucose, fructose, ribulose and glyceric acid, as well as the compounds of the Hatch-Slack pathway that include oxaloacetate, malate and pyruvate. Thus, supplying the heterotrophically cultured plant cells with one of the before-mentioned carbon sources and the other required minimal nutrients of the medium results in growth and proliferation of the plant cells in that the plant cells contain endogenous enzymes that utilize such compounds for growth and proliferation.

Inasmuch as living organisms all utilize carbon sources for growth and proliferation, many such enzymes are available for use. An enzyme that converts a non-useful carbon source to a useful 6-carbon source such as glucose or fructose is preferred, and such enzymes are utilized illustratively herein.

One preferred enzyme used illustratively herein is phosphomannose isomerase (*pmi/mana*; E.C. 5.3.1.8) that converts mannose-6-phosphate into fructose-6-phosphate; the latter compound that supports plant cell growth and proliferation. When mannose is used as the encrypted carbon source, mannose-6-phosphate is formed in the cell by the enzymatic action of an endogenous kinase. The phosphomannose isomerase gene is designated as the *pmi* gene in *Rhizobium meliloti* (*Gene*, 122: 35-43), *Pseudomonas aeruginosa* (*Gene*, 42:293-302), *Saccharomyces cerevisiae* (*Molecular and Cellular Biology*, 12:2924-2930) and *Salmonella typhimurium* (*Gene*, 103:135-136). In *Escherichia coli*, the phosphomannose isomerase gene is called *mana* (*Gene*, 32:41-48).

Another useful enzyme is L-idoitol dehydrogenase (EC 1.1.1.14) that converts sorbitol into fructose. Several aldehydoreductase enzymes are known that can be used to convert sorbitol as encrypted carbon source into the useful carbon source glucose. One exemplary enzyme is D-sorbitol 1-oxidoreductase (EC 1.1.00.24). Lactase (EC 3.2.1.108) that converts the encrypted carbon source lactose into glucose and galactose, as do several other β -galactosidases, and α , α -trehalase (EC 3.2.1.28) that converts the encrypted carbon source α , α -trehalose into glucose are yet other useful enzymes.

Upon determination of the heterologous marker enzyme for use, the gene that encodes that enzyme is obtained. The genes that encode many heterologous marker enzymes have been reported in the literature and can be obtained from the authors, as was the case of the *pmi* gene used here. Many useful gene sequences are also reported in the GenBank database such as the human L-idoitol dehydrogenase gene (GenBank accession number L29008) or the rat aldehyde reductase genes (GenBank accession numbers X74673 and D10484) or the human aldehyde reductase gene (GenBank accession number J04794). Sequences for several useful β -galactosidases are also available in the GenBank data base. For example, the enzyme from *Brassica oleracea* has accession Number X84684, that of the Granny Smith apple (*Malus domestica*) has accession Number L29451, and three genes from *E. coli* have the accession Numbers X03228, M13700 and M13797. The sequence of the *S.typhimurium pmi* gene used illustratively here has GenBank accession Number X57117.

Where the desired gene is unavailable, but its sequence is known, PCR technology can be used with a DNA or RNA prepared from a source reported or otherwise known to contain the gene to obtain a useful copy of that gene.

Where a sequence of a desired gene is not available, the enzyme itself is obtained following usual procedures, and its termini are sequenced sufficiently so that sets of redundant DNA probes can be prepared and used with PCR technology to obtain a DNA copy of the gene from a DNA or RNA library of the organism from which the enzyme was isolated.

The preparation and use of DNA and RNA libraries for obtaining desired genes from host organisms is well known in this art and will not be dealt with further herein.

The DNA segment encoding the heterologous selectable marker enzyme (first gene) is typically isolated and purified containing appropriate endonuclease restriction sites at the 5'- and 3'-termini so that that gene can be operatively linked to the promoter and termination DNA segments. If that DNA segment is not so obtained, suitable restriction sites can be added by *in vitro* ligation or by techniques that are well known.

An above-described first gene is operatively linked to a promoter DNA segment. That promoter is a segment that is operative in plant cells. Exemplary useful promoters include the constitutive CaMV 35S promoter of the cauliflower mosaic virus, the octopine synthase promoter (P-Ocs) and the nopaline synthase promoter (P-Nos) that are constitutive promoters.

In one embodiment, the promoter utilized is repressed by a product of the normal metabolism of the transgenic (or non-transgenic) plant cultured under autotrophic growth conditions. Thus, once a mature plant has been regenerated, and selection is no longer necessary, the promoter controlling expression of the marker enzyme is shut off or repressed so that none or a reduced amount of the selecting enzyme is expressed.

An exemplary promoter whose activity is repressed by a normal product of plant metabolism is

the rice α -amylase Amy3A promoter [Thomas et al., Plant Physiol., 106:1235-1239(1994)]. Expression from this α -amylase promoter in rice is repressed by the presence of sucrose, a product of the normal, autotrophic, metabolism of both transformed and non-transformed rice cells.

Two additional promoters that are repressible by normal products of plant metabolism are the cucumber malate synthase and isocitrate lyase promoters reported in Graham et al., Plant Cell, 6: 761-772 (1994). Glucose, fructose and raffinose in culture media repressed both promoters. Inasmuch as the malate synthase promoter is repressed when plant cells are in the presence of mannose, that promoter is not utilized in conjunction with the *pmi/mana* gene and mannose as the first encrypted carbon source.

The final element of the first cassette is a first termination DNA segment that is operatively linked to the 3'-end of the first gene. Several termination segments useful in plants are well known and can be used herein. One exemplary segment is the 3'-non-translated region of the nopaline synthase gene [Nos-T; Fraley et al., Proc. Natl. Acad. Sci. USA, 80:4803-4807(1983)] used herein. Another is the 3'-non-translated region of the pea *rbcS-E9* gene [E9; Coruzzi et al., Embo J., 3:1671-1679(1984)].

The promoter and termination DNA segments are also preferably terminated by appropriate endonuclease restriction sites for ligation and operable linkage to the termini of the first gene and ligation to the second expression cassette as part of the genomic heterologous DNA segment, and also into an appropriate vector. Blunt end ligation can also be used for operative linking to the first gene and/or a vector.

The second expression cassette contains a second gene that is expressed in a transformed plant. That second or target gene encodes for the expression of a protein in a transformed plant.

-23-

The second gene can be any gene that is desired. Exemplary genes include those listed in Table I, below, whose transformations into plants have been disclosed in the patent citations shown in that Table.

5

TABLE 1

	<u>Second Gene Product</u>	<u>Citation</u>
	Antibody	U.S. 5,202,422
10	Mammalian Peptide	WO 87/00865
	HMG-CoA Reductase	U.S. 5,306,862
	Phosphofructokinase	U.S. 5,387,756
15	Waxy Locus of Wheat (Antisense)	U.S. 5,365,016
20	ADP-Glucose pyrophosphorylase, antisense	EP 0 368 506 A2 EP 0 455 316 A2 WO 92/11382
	Potato L-amylase	EP 0 470 145 B1
25	Sucrose phosphate Synthase	EP 0 466 995 A2 EP 0 530 978
	<i>E.coli</i> inorganic pyrophosphorylase	EP 0 485 044 A2
30	Maize 1,4- α -glucan branching enzyme, antisense	WO 92/11375
35	Granule-bound starch synthase, antisense	WO 92/11376
	Tomato vacuolar	WO 92/14831

invertase, antisense

E. Herbicola genes
in the carotenoid
synthetic pathway

WO 91/13078

5
10 The second gene is operatively linked to a second promoter DNA segment that controls expression of the second gene. This promoter is not repressed by a product of normal plant metabolism, and can be a constitutive promoter such as the CaMV 35S, P-Ocs and P-Nos promoters discussed before, or an organ-enhanced promoter that causes expression in one or more limited organs of the transformed plant.

15 Expression in one or more preselected organs with little or no expression in other organs such as roots versus leaves or stems is referred to herein as enhanced or preferential expression.

20 The final element in the second expression cassette is a termination DNA segment. Any termination segment can be used, as was discussed before in regard to the first cassette.

25 The elements of the second expression cassette are operatively linked together, usually via ligated endonuclease sites, although blunt end ligation can also be utilized, as before.

30 It is also noted that either or both of the heterologous enzyme encoded by the first gene and the product of the second gene can be portions of a polyprotein that actually contains two or more proteins, e.g. enzymes, linked together by a peptide bond as has been found for several viral genes. See, for example, Vardi et al., Proc. Natl. Acad. Sci. USA, 90:7413-7417(1993) and Maiti et al., Proc. Natl. Acad. USA, 90:6110-6114(1993).

35 The first and second expression cassettes are linked together in the genomic heterologous DNA segment so that the two expression cassettes constitute very

tightly linked loci that are transformed into the plant's chromosomal DNA. Thus, as is the case with other selectable marker systems, transformation of plant cells with one of the cassettes is very strongly correlated with transformation with the other. In this instance, growth and proliferation of cells that were putatively transformed indicates that the transformation with the second cassette was successful.

Use of the DNA segment containing the two linked expression cassettes to transform plant cells is discussed in detail hereinafter.

In some instances, it is desirable to transform plant cells on two separate occasions. Such double transformations require the sequential use of two selectable markers and two target genes. In the terminology used herein, such a double transformation utilizes four expression cassettes and two encrypted (growth-limiting) sources of carbon for heterotrophic growth. Two heterologous genomic DNA segments are present in the twice-transformed plants.

For ease of discussion as to this embodiment, the first and second expression cassettes are those already discussed, and those expression cassettes are present in the first-named genomic heterologous DNA segments. A second genomic heterologous DNA segment is also present in a twice-transformed plant and contains a third and a fourth expression cassettes.

The third expression cassette contains a second heterologous DNA selectable marker segment that includes (i) a second heterologous gene that encodes a second heterologous enzyme that is operatively linked to (ii) a third promoter DNA sequence that controls the expression of the second heterologous gene and (iii) a termination DNA segment.

The third promoter DNA can be the same promoter used with the first gene that encodes a heterologous enzyme. The termination DNA segment is also any termination segment useful in plants as is

also discussed before. The third promoter and this termination DNA segment are linked as discussed previously.

5 The second gene that encodes a second heterologous enzyme is different from those genes discussed before. The second heterologous enzyme converts a second encrypted carbon source into the first-named encrypted carbon source that supports growth and proliferation of the transformed plant
10 cells. That second encrypted carbon source also does not support growth and proliferation of non-transformed plant cells of the same type.

 Thus, the second encrypted carbon source can be viewed as a precursor to the first-named encrypted
15 carbon source that is converted into a useful source of carbon to support cell growth and proliferation by the first heterologous gene. More specifically, where mannose is illustratively the first encrypted carbon source and phosphomannose isomerase is the first
20 heterologous gene, mannitol can be used as the second encrypted carbon source along with mannitol 1-oxidoreductase as the second heterologous gene.

 Thus, plant cells transformed with a first and second expression cassette containing the *pmi/manA*
25 gene as a first gene selectable marker are transformed again with third and fourth expression cassettes whose third expression cassette contains a gene that encodes the mannitol 1-oxidoreductase gene and whose fourth cassette contains another desired gene. The resulting
30 mixture of twice-transformed and once-transformed plant cells is heterotrophically cultured in a minimal nutrient medium that contains mannitol as the carbon source. Inasmuch as only the twice transformed cells contain a gene encoding mannitol 1-oxidoreductase and
35 phosphomannose isomerase, only those cells grow and proliferate because they convert mannitol to mannose and mannose into fructose-6-phosphate.

Those proliferating twice-transformed cells are preferably recovered and regenerated into mature plants as discussed elsewhere.

It is noted that reference herein to "first", "second", "third" and "fourth" is for convenience only. Thus, the second cassette can be upstream of the first cassette, as can the fourth cassette be upstream of the third cassette.

In a second embodiment, a process for selectively increasing the number of transformed plant cells regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions is contemplated. In accordance with this process,

(a) a mixture of transformed and non-transformed plant cells is cultured under heterotrophic culture conditions in a culture medium that contains standard nutrients required for proliferation and differentiation by those plant cells except for a useful source of carbon and 1.5 to 3 times the standard amount of phosphorus. The normally present useful source of carbon is replaced by an encrypted, latent or growth-limiting carbon source that does not support growth and proliferation of the non-transformed cells. The transformed cells of the mixture contain a genomic (chromosomally integrated) heterologous DNA segment that contains at least one expression cassette.

At least one expression cassette contains a heterologous DNA selectable marker segment that includes (i) a heterologous gene that encodes a heterologous enzyme that on expression converts the encrypted carbon source into a useful carbon source that supports growth and proliferation of the transformed plant cells under heterotrophic culture conditions. The gene is operatively linked to (ii) a first promoter DNA segment that controls expression of the heterologous gene, and (iii) a termination DNA segment. This expression cassette may be identical to

the first expression cassette described earlier as the first embodiment of this invention. Therefore, the previous descriptions relating to the first expression cassette may be applicable to the expression cassette
5 used in the second embodiment of this invention.

The heterologous DNA segment must contain at least the expression cassette described above. Optionally, the heterologous DNA segment may contain additional expression cassettes which contain any
10 gene(s) that are expressed in transformed plants. These expression cassettes may be identical to the second through fourth expression cassettes described above with respect to the first embodiment of this invention. Therefore, the previous descriptions
15 relating to these expression cassettes may be applicable in the second embodiment of this invention.

(b) The heterotrophic culture conditions are maintained for a time period sufficient for the transformed plant cells to express the heterologous
20 enzyme, proliferate and grow.

The second embodiment of the invention utilizes the discovery that when additional phosphorous is added to the culture medium containing the minimal nutrients required for growth and proliferation of the
25 non-transformed plant cells except for an encrypted or latent carbon source that does not support growth and proliferation of the non-transformed plant cells, that there is an increase in the quantity and quality of transformed plant cells regenerated.

"The standard amount of phosphorous" as used herein refers to the amount or level of phosphorous contained in the commonly used Murashige and Skoog culture medium which is frequently used in this art. The Murashige and Skoog culture medium typically
30 contains 170.00 mg/L of phosphorous such as potassium phosphate monobasic. Nonetheless, it is well known in the art that the Murashige and Skoog culture medium can be modified to meet certain requirements. Generally,

when the Murashige and Skoog medium is modified, these modifications are in the composition of microelements, vitamins or inclusion of other additives. An example of a modified Murashige and Skoog medium in which only the vitamins are altered is the Linsmaier and Skoog medium (described in Physiologia Plantarum 18: 100-127 (1965)). This medium contains 170.00 mg/L of phosphorous.

In this invention, the inventors have discovered that when the culture medium that contains the encrypted or latent carbon source also contains 1.5 to about 3 times the standard amount of phosphorous normally contained in said medium that an enhanced number of regenerated transformed plant cells can be obtained. Additionally, among those regenerates, a higher percentage of normal shoots can be recovered as well.

While not wishing to be bound by any theory, the inventors believe that the reason why transformation experiments on the growth medium containing the encrypted or latent carbon source with supplemented levels of phosphate are more successful than experiments on the growth medium containing only the encrypted carbon source is because that when the growth medium contains the additional levels of phosphate the transformed plant cells have at least two growth advantages over the non-transformed plant cells. The first advantage is that the transformed plant cells can access additional carbon from the medium. The second advantage is that the phosphate confers an additional growth advantage to transformed cells capable of growing on the limited source of carbon over the non-transformed plant cells which are not capable of growing on such a limited carbon source.

The phosphorous added to the culture medium can be any water soluble phosphate salt. For example, potassium phosphate (monobasic, dibasic and tribasic), sodium phosphate (monobasic, and dibasic) calcium

-30-

phosphate (monobasic, dibasic and tribasic), sodium hexameta-phosphate, ammonium phosphate (monobasic and dibasic), orthophosphoric acid, ferric phosphate, sodium glycerophosphate, ferric glycerophosphate and
5 organic phosphorous (such as ATP, ADP, etc) can be used.

In a third embodiment, a process for selectively increasing the number of transformed plant cells regenerated from a mixture of transformed and
10 non-transformed plant cells cultured under delayed heterotrophic culture conditions is contemplated. In accordance with this process,

(a) a mixture of transformed and non-transformed plant cells is cultured for up to two weeks
15 in a first culture medium that contains the minimal nutrients required for proliferation and growth by non-transformed plant cells including a source of carbon that supports growth and proliferation. The transformed plants cells containing a genomic heterologous DNA
20 segment that contains at least one expression cassette.

At least one expression cassette contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts an
25 encrypted carbon source into a carbon source that supports growth and proliferation of said transformed plant cells under heterotrophic culture conditions, said first gene being operatively linked to (ii) a first promoter DNA segment that controls expression of
30 said heterologous gene, and (iii) a termination DNA segment. This expression cassette may be identical to the first expression cassette described earlier as the first embodiment of this invention. Therefore, the previous descriptions relating to the first expression
35 cassette in the first embodiment of this invention may be applicable to the expression cassette used in the third embodiment of this invention.

The heterologous DNA segment must contain at least the expression cassette described above. Optionally, the heterologous DNA segment may contain additional expression cassettes which contain any gene(s) that are expressed in transformed plants. Therefore, the previous description in the first embodiment relating to such expression cassettes may be applicable to the third embodiment of this invention.

(b) After sufficient time in the first medium, transformed and non-transformed plant cells are removed from the first culture medium.

(c) After removal from the first culture medium, the transformed and non-transformed plant cells are placed under heterotrophic culture conditions in a second culture medium that contains the minimal nutrients required for proliferation and growth of the non-transformed plant cells except for an encrypted carbon source that does not support growth and proliferation of said non-transformed plant cells and 1.5 to 3 times the standard amount of phosphorous.

(d) The heterotrophic culture conditions are maintained for a time period sufficient for said transformed plant cells to express said heterologous enzyme, proliferate and grow.

As discussed earlier with respect to the second embodiment, the addition of 1.5 to 3 times the standard amount of phosphorous increases the quantity and quality of transformed plant cells regenerated. Also as in the second embodiment, the phosphorous added to the culture medium can be any water soluble phosphate salt.

Recombinant DNA Molecules

A recombinant DNA molecule useful herein can be produced by operatively linking a vector to an isolated heterologous DNA segment that contains two expression cassettes to form a plasmid. Particularly preferred recombinant DNA molecules are discussed in detail in the examples, hereafter. Vectors capable of

directing the expression of the gene are referred to herein as "expression vectors".

5 In one preferred embodiment, a vector includes a prokaryotic replicon; i.e., a DNA sequence having the ability to direct autonomous replication and maintenance of the recombinant DNA molecule extrachromosomally in a prokaryotic host cell transformed therewith. Such replicons are well known in the art.

10 Those vectors that include a prokaryotic replicon can also include a prokaryotic promoter region capable of directing the expression of gene in a host cell, such as *E. coli*, transformed therewith. Promoter sequences compatible with bacterial hosts are typically provided in plasmid vectors containing one or more
15 convenient restriction sites for insertion of a DNA segment of the present invention. Typical of such vector plasmids are pUC18, pUC19, and pBR322 available from Gibco BRL, Gaithersburg, MD. These vectors are
20 utilized in the synthesis of the DNA segments present in the integrating expression vectors.

Typical vectors useful for expression of genes in higher plants are well known in the art and include vectors that contain portions of the tumor-inducing (Ti) plasmid of *Agrobacterium tumefaciens* described by Rogers et al., Meth. in Enzymol., 153:253-277 (1987). These vectors are plant integrating
25 vectors in that on transformation, a portion of the vector DNA is integrated into the genome of the host plant. For integrating vectors containing portions of the Ti plasmid, the region integrated into the host plant chromosomes is that between the right and left borders of the Ti plasmid, or TDNA.
30

A preferred plant transformation vector useful herein is vector BIN19 [Bevan, Nucleic Acids Res., 12:8711-8721(1984)]. Other exemplary *A. tumefaciens* vectors useful herein are plasmids pKYLX6 and pKYLX7 of Schardl et al., Gene, 61:1-11 (1987) and
35

Berger et al., Proc. Natl. Acad. Sci. USA, 86:8402-8406 (1989). Another useful vector herein is plasmid pBI101 that is available from Clontech Laboratories, Inc., Palo Alto, CA. Plasmids BIN19, pKYLX7, pKYLX71 and
5 pBI101 are binary vectors that are used in *A. tumefaciens*.

Another plant transformation system is based on *Agrobacterium rhizogenes* that induces hairy roots rather than a tumor on transformation. Application
10 PCT/US87/02512 (WO 88/02405 published April 7, 1988) describes the use of *A. rhizogenes* strain A4 and its Ri plasmid along with *A. tumefaciens* vectors pARC8 or pARC16 to transform the cucumber *Cucumis sativus*, cv, Straight Eight, and form regenerated cucumber plants.

15 An *Agrobacterium*-based transformation system for melon (*Cucumis melo*), which is another member of the cucumber family of *Cucurbitaceae*, was reported by Dong et al., Bio/Technology, 9:858-863 (1991). Those workers used a binary vector that utilized the
20 constitutive CaMV 35S promoter, and found evidence of transformation via a reporter gene in substantially all tissues examined. That work illustrates the amenability of melons to transformation via *A. tumefaciens*.

25 A variety of methods has been developed to operatively link DNA segments into vectors via complementary cohesive termini or blunt ends. In addition, synthetic linkers containing a variety of restriction endonuclease sites are commercially
30 available from a number of sources including New England BioLabs, Beverly, MA. These methods and use of synthetic linkers are well know in the art and will not be discussed further here. See Sambrook et al., Molecular Cloning, 2ed; Cold Spring Harbor Laboratory
35 Press (1989).

Introducing genes into higher plants

Methods for transforming higher, multicellular flowering plants include *Agrobacterium*-mediated plant cell transformation, protoplast transformation, gene transfer into pollen, injection
5 into reproductive organs, transformation of microspores and injection into immature embryos. Each of these methods has distinct advantages and disadvantages. Thus, one particular method of introducing genes into a particular plant species may not necessarily be the
10 most effective for another plant species, but it is well known which methods are useful for a particular plant species.

Agrobacterium-mediated transfer is a widely applicable system for introducing genes into plant
15 cells. The use of *Agrobacterium*-mediated expression vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described by Fraley et al., Biotechnology, 3:629 (1985) and Rogers et al., Methods in Enzymology, 153:253-277 (1987).
20 Further, the integration of the T-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is within the border sequences, and intervening DNA is usually inserted into the plant genome as described by Spielmann et al., Mol.
25 Gen. Genet., 205:34 (1986) and Jorgensen et al., Mol. Gen. Genet., 207:471 (1987).

Agrobacterium transformation vectors such as those discussed before are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient
30 manipulations as described by Klee et al., in Plant DNA Infectious Agents, T. Hohn and J. Schell, eds., Springer-Verlag, New York (1985) pp. 179-203.

Moreover, technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved
35 the arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. The vectors described by Rogers et al., Methods in

Enzymology, 153:253 (1987), have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes.

In those plant species where *Agrobacterium*-mediated transformation is efficient; i.e., in dicotyledonous plants, it is a method of choice because of the facile and defined nature of the gene transfer. Particularly preferred dicotyledonous plants include those of *Brassicaceae* (cabbage and broccoli), *Compositae* (lettuce), *Umbelliferae* (carrots), *Solanaceae* (pepper and tomato) and *Cucurbitaceae* (melon and squash) families of higher plants.

Monocots such as cereals and grasses appear not to be natural hosts for *Agrobacterium*, although transgenic plants have been produced in asparagus using *Agrobacterium* vectors as described by Bytebier et al., Proc. Natl. Acad. Sci. U.S.A., 84:5345 (1987). Commercially important cereal grains such as rice, corn, oats and wheat are usually transformed using alternative methods.

To transform plant species that cannot be successfully transformed using *Agrobacterium*, a "particle gun" or high-velocity microprojectile technology can be utilized. Using such technology, DNA is carried through the cell wall and into the cytoplasm on the surface of small metal particles as described in Klein et al., Nature, 327:70 (1987); Klein et al., Proc. Natl. Acad. Sci. U.S.A., 85:8502 (1988); and McCabe et al., Biotechnology, 6:923 (1988); and Vasil et al., Bio/Technology, 9:667-674 (1992). The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants.

The microprojectile mediated transformation is a preferred source of transgenic cells from monocotyledonous plants. Particularly preferred

monocotyledonous plants include maize, wheat, oats, barley, rice, sorghum and forage grasses. Metal particles coated with DNA have been used to successfully transform maize and wheat cells. Fertile, stably transformed tobacco and soybean plants have also been transformed by this method. Regeneration of cereals from immature embryos or explants can be effected as described by Vasil, Biotechnology, 6:397 (1988).

Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation, and combinations of these treatments. See, for example, Potrykus et al., Mol. Gen. Genet., 199:183 (1985); Lorz et al., Mol. Gen. Genet., 199:178 (1985); Fromm et al., Nature, 319:791 (1986); Uchimiya et al., Mol. Gen. Genet., 204:204 (1986); Callis et al., Genes and Development, 1:1183 (1987); Marcotte et al., Nature, 335:454 (1988); Wang et al., Bio/Technology, 10:691-696 (1992); and Fennell et al., Plant Cell Reports, 11:567-570 (1992).

Application of these systems to different plant species depends upon the ability to regenerate that particular plant species from protoplasts. Illustrative methods for the regeneration of cereals from protoplasts are described in Fujimura et al., Plant Tissue Culture Letters, 2:74 (1985); Toriyama et al., Theor. Appl. Genet., 73:16 (1986); Yamada et al., Plant Cell Rep., 4:85 (1986); Abdullah et al., Biotechnology, 4:1087 (1986). U.S. Patent No. 4,634,674 teaches regeneration of tomato plants from protoplasts.

DNA can also be introduced into plants by direct DNA transfer into pollen as described by Zhou et al., Methods in Enzymology, 101:433 (1983); D. Hess, Intern. Rev. Cytol., 107:367 (1987); Luo et al., Plant Mol. Biol. Reporter, 6:165 (1988). Expression of transformed genes can be obtained following injection

of the DNA into reproductive organs of a plant as described by Pena et al., Nature, 325:274 (1987). U.S. Patent No. 5,302,523 teaches transformation of maize cells using whisker bodies to penetrate the cells. DNA can also be injected directly into the cells of immature embryos and during the rehydration of desiccated embryos as described by Neuhaus et al., Theor. Appl. Genet., 75:30 (1987); and Bedbrook et al., in Proceedings Bio Expo 1986, Butterworth, Stoneham, MA, pp. 27-54 (1986), as well as by poly(ethylene glycol) or electroporation into maize microspores from which dihaploid homozygous plants can be regenerated (Fennell et al., Plant Cell Reports, 11:567-570 (1992)).

The regeneration of plants from transformed plant cells such as either single plant protoplasts or various explants is well known in the art. See, for example, Methods for Plant Molecular Biology, A. Weissbach and H. Weissbach, eds., Academic Press, Inc., San Diego, CA (1988). The regeneration and growth process includes the steps of selection of transformant cells and shoots, rooting the transformant shoots and growth of the plantlets in soil.

Growth media for plant cells contemplated herein are those familiar to workers of ordinary skill in these arts except that an encrypted carbon source replaces the usually used carbon source such as glucose or sucrose. Exemplary media contain Murashige and Skoog (MS) salts and RO vitamins. Regeneration media are similarly well known and also contain plant hormones such as benzyladenine, gibberellic acid, indoleacetic acid, naphthylacetic acid, zeatine, thiamine hydrochloride and the like, and utilize an encrypted carbon source.

As discussed earlier, one special feature of a preferred nutrient medium for regeneration is that from approximately 1.5 to about 3 times the standard amount of phosphorous is present as a water-soluble

phosphate salt. The basis for this preference is the finding that an enhanced number of healthy regenerated shoots were obtained using the enhanced amount of phosphate salt as well as a greater percentage of normal shoots among those regenerates.

The regeneration of plants containing the foreign gene introduced by *Agrobacterium* from leaf explants can be achieved as described by Horsch et al., Science, 227:1229-1231 (1985). In this procedure, transformants are grown in the presence of the appropriate encrypted carbon source in a medium that induces the regeneration of shoots in the plant species being transformed as described by Fraley et al., Proc. Natl. Acad. Sci. U.S.A., 80:4803 (1983). This procedure typically produces shoots within two to four months and these transformant shoots are then transferred to an appropriate root-inducing medium and an antibiotic to prevent bacterial growth.

Transformant shoots that begin to root in the presence of an encrypted carbon source and form plantlets are then transplanted to soil or other media to allow the production of roots. These procedures vary depending upon the particular plant species employed, such variations being well known in the art.

Genetics

A transgenic plant formed using *Agrobacterium* transformation methods can contain a single transgene on one chromosome. Such transgenic plants can be referred to as being heterozygous for the added gene. However, inasmuch as use of the word "heterozygous" usually implies the presence of a complementary gene at the same locus of the second chromosome of a pair of chromosomes, and there is no such gene in a plant containing one added gene as here, a more accurate name for such a plant is a hemizygote. The original transgenic plant formed using *Agrobacterium* is referred to as a hemizygote.

More preferred is a transgenic plant that is homozygous for the added expression cassettes, e.g., a transgenic plant that contains two sets of added expression cassettes, one set at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can exhibit at least enhanced expression of the selectable marker as compared to both a native, non-transgenic plant and an independent segregant transgenic plant. A homozygous transgenic plant can also be prepared from transformed microspores, as noted before. Even more preferred is a plant created by a cross of different inbred lines. Plant seed produced by such a cross are referred to as hybrid seed. There are many benefits to growing hybrid seed that include, a more uniform product, higher yields, and healthier plants. Collectively, these advantages are well known in agriculture and are often referred to as hybrid vigor. If one of the inbred parents carry the transgenic cassettes in a homozygous condition, then the hybrid progeny would all contain a single copy of the transgenic cassettes (i.e., they would be hemizygotic for the transgenes). In such a crossing scheme, the hybrid progeny will have both the advantages of hybrid vigor and the advantage of the effects conferred by the transgenes.

It is to be understood that two different transgenic plants can be mated to produce offspring that contain two sets of independently added segregating cassette sets. Selfing of appropriate progeny can produce plants that are homozygous for both added cassette sets. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated.

A transgenic plant of this invention thus has heterologous DNA containing two expression cassettes (a set). A preferred transgenic plant is an independent segregant for the added set of cassettes and can transmit those cassettes and their activity to its

progeny. A more preferred transgenic plant is homozygous for those heterologous cassette sets, and transmits those genes to all of its offspring on sexual mating.

5

Plant Transformation Kit

Another aspect of this invention is a kit for transforming plant cells. This kit, which itself is usually contained in a package, contains two packages and also preferably contains a set of directions for use of the kit.

A first package contains a DNA segment heterologous to the plant to be transformed. That DNA segment contains an expression cassette operatively linked to a linker segment that includes at least one restriction endonuclease site, and preferably includes a plurality of such sites. The expression cassette is a first (or third) expression cassette discussed before that contains a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that converts an encrypted carbon source that does not support growth and proliferation of non-transformed plant cells into a carbon source that supports growth and proliferation of the transformed cells upon expression during heterotrophic culture of the transformed plant cells. That gene is operatively linked to (ii) a promoter DNA segment that controls the expression of that heterologous gene and (iii) a termination DNA segment.

The first gene can encode any of the before-discussed heterologous genes, with the *pmi* gene being one preferred gene. Another preferred gene is the mannitol 1-oxidoreductase gene that converts mannitol into mannose and is useful for situations where double transformation is contemplated.

The promoter and termination DNA segments are those discussed before in connection with the first

expression cassette. A repressible promoter DNA segment is particularly preferred.

5 The linker segment of this DNA preferably contains a plurality of endonuclease restriction sites as are commonly found in vectors. This linker and its one or more restriction sites facilitates incorporation of a second expression cassette of the user's choice into the heterologous DNA segment. The linker segment can be located upstream or downstream of the first
10 expression cassette.

The DNA of the first package is itself preferably within the TDNA borders of the Ti plasmid of *Agrobacterium tumefaciens* in a vector adapted for *Agrobacterium*-mediated plant transformation. Thus, an
15 expression vector is contemplated for *Agrobacterium*-mediated plant transformation that contains all of the elements needed for that transformation except for the second expression cassette containing the second, target, gene and its promoter and termination segments
20 that are supplied by the user.

Another embodiment of this aspect of the invention includes one or the other or both of 5'-promoter and 3'-termination DNA segments with an operatively-linked linker containing at least one
25 restriction site in between so that all that the user need do is insert (ligate) a desired second, target, gene. These segments can be located together upstream or downstream of the recited expression cassette.

The second package of the kit contains
30 minimal nutrients required for growth and proliferation of non-transformed plant cells during heterotrophic culture, except for a source of carbon. The source of carbon usually present in such nutrient media is replaced by an encrypted carbon source that does not
35 support growth and proliferation of non-transformed plant cells but supports growth and proliferation of transformed plant cells whose genome contains a before-described DNA segment containing a first expression

cassette.

5 A nutrient medium contemplated here as present in the second package can be any useful medium such as any of the media discussed herein like MS medium containing MS salts and RO vitamins that is free of a usual carbon source. The encrypted carbon source is also a previously discussed encrypted carbon source such as mannose, lactose, sorbitol or mannitol as a second encrypted carbon source.

10 A third package can also be included that contains components of a regeneration medium and the same encrypted carbon source. Thus, nutrient materials such as MS salts and RO vitamins are included along with plant hormones such as benzyladenine and
15 gibberellic acid, and the like as are well known. Antibiotics such as carbenicillin and timentin or ticarcillin that eliminate *Agrobacteria* can also be included. Each of the ingredients is present in its usually-used amount except for phosphorous that is
20 preferably present at about 1.5 to about 3 times its usual amount in such a medium as a water-soluble phosphate salt.

Results

25 The first step in the heterotrophic complementation selection system is to identify a carbon source such as a carbohydrate that is not metabolized by plant cells to support growth and proliferation. Examples of such carbohydrates are
30 mannose, mannitol, sorbitol, lactose, trehalose and salicin.

The second step in the carbon source-based selection system is to select a gene that encodes an enzyme that permits the transformed tissue to utilize
35 that encrypted carbon source, whereas that carbon source is not normally metabolized by non-transformed plant tissue. One particular exemplary gene encodes the enzyme phosphomannose isomerase (*pmi*; EC 5.3.1.8).

This enzyme reversibly catalyzes the conversion of fructose 6-phosphate to mannose 6-phosphate. This gene is designated as the *pmi* gene in *Rhizobium meliloti* (Gene, 122: 35-43), *Pseudomonas aeruginosa* (Gene, 42:293-302), *Saccharomyces cerevisiae* (Molecular and Cellular Biology, 12:2924-2930) and *Salmonella typhimurium* (Gene, 103:135-136). In *Escherichia coli*, the phosphomannose isomerase gene is called *manA* (Gene, 32:41-48).

Selection of a Growth-limiting Carbohydrate

All media used herein was that of Murashige and Skoog, Physiol. Plant, 15:437-498 (1962) (Medium MS) salts and RO vitamins (composition listed below) that were adjusted to pH = 5.7 and solidified with 9 g/L of Noble Agar (Gibco). Medium R1F was MS supplemented with 1 mg/L indoleacetic acid (IAA), 0.65 mg/L zeatine and 16 g/L glucose. Medium MR1 was MS supplemented with 1 mg/L IAA, 0.65 mg/L zeatine and 16 g/L mannose. Medium MR1/2 was MS supplemented with 0.5 mg/L IAA, 0.325 mg/L of zeatine and 16 g/L mannose. Medium RO was MS supplemented with 16 g/L glucose. Medium MRO was MS supplemented with 16 g/L mannose.

- 44 -

Table 1.

Composition of media used in tomato regeneration.

	<u>Medium MS Salts</u>	<u>mg/L</u>
5	Ammonium nitrate	1650.000
	Boric acid	6.200
	Calcium chloride	440.000
	Cobaltous chloride	0.025
	Cupric sulfate pentahydrate	0.025
10	Ferrous sulfate septahydrate	27.800
	Magnesium sulfate septahydrate	370.000
	Manganese sulfate monohydrate	15.600
	Potassium iodide	0.083
15	Potassium nitrate	1900.000
	Potassium phosphate monobasic	170.000
	Sodium ethylenediamine tetraacetate	
	Sodium molybdate dihydrate	0.250
	Zinc sulfate septahydrate	8.600
20	<u>RO Vitamins</u>	<u>mg/L</u>
	Nicotinic acid	5.000
	Thiamine HCl	0.500
	Pyridoxine	0.500
	Myo-inositol	100.000
25	Glycine	

Tomato seeds were sterilized in a solution containing 20 percent of a commercial household bleach containing 5.25 percent sodium hypochlorite for 20 minutes, rinsed 3 times in sterile distilled water and

30 planted on RO medium solidified with 10 g/L of Noble agar (Gibco) in 135 mm Phytakon™ tissue culture vessels (Sigma, St. Louis, MO). Seeds were germinated for 72

hours at 25°C in the dark, then moved to a lighted shelf under approximately $80 \mu\text{molm}^{-2}\text{s}^{-1}$ photosynthetic photon flux density (PPFD), at 24-26°C. Plant tissue used for regeneration was prepared by removing cotyledons from 7-day-old seedlings and cutting them into three parts (proximal, middle and distal to the growing point). The middle and proximal parts were used for the evaluation of growth on potential non-useful carbohydrates.

The following carbohydrates were considered for use as a non-useful encrypted, replacement of a carbon source in plant regeneration medium: mannose, mannitol, sorbitol, lactose, α,α -trehalose and salicin. Different R1F media were prepared, each containing one of those carbohydrates added at 0.09 M concentration as a sole carbon source. Control R1F medium contained 0.09M (16 g/L) glucose as a carbon source. Tomato tissue was placed abaxial side down on the media and incubated in sealed petri plates at 24-26 °C, under $30 \mu\text{molm}^{-2}\text{s}^{-1}$ PPFD, 16 hours photoperiod. Tomato tissue was observed for signs of regeneration weekly.

Throughout the carbohydrate evaluation, which lasted four weeks, none of the altered media containing the above-listed carbohydrates supported regeneration of tomato shoots, whereas tomato tissue placed on the control glucose-supplemented R1F medium produced abundant shoots.

Selection of a Gene to Convert an Encrypted Carbohydrate

It is known that plants can take mannose up from media, and subsequently phosphorylate mannose into mannose-6-phosphate. Sheu-Hwa et al. New Phyto., 74:383-392 (1975). It is concluded that mannose-6-phosphate does not support growth and proliferation of plant cells because plant cells cultured using mannose as the sole carbon source do not grow and proliferate.

A gene was therefore chosen that encodes an enzyme that converts mannose-6-phosphate into fructose-6-phosphate. Fructose-6-phosphate is a compound known to be metabolized by plants. Thus, plant cells that express this gene are able to utilize mannose to grow and proliferate, and have a selective growth advantage when placed on media containing mannose as the sole carbon source.

Construction of the Transfer Vector

10 A *pmi* gene from a prokaryotic species was selected and used to create a plant transformation vector that included a selectable marker of the present invention. Molecular biology techniques, that are well known to those skilled in the art, were used to create
15 the plant transformation vector containing the selectable marker gene. The steps used to move the *pmi* gene into a plant transformation vector are shown in Fig. 2. Enzymes used to manipulate DNA can be purchased either from New England Biolabs or Boehringer Mannheim.
20

Briefly, plasmid pADE253, containing the *S. typhimurium pmi* gene (GenBank accession number X57117), was a gift from Dr. James Hackett, University of Adelaide. Plasmid pADE253 is a derivative of the
25 commonly used *E. coli* cloning vector pBR322 that contains a 1650 base pair (bp) *Cla*I fragment (Fig. 2) that contains the *pmi* gene. SEQ ID NO:1 shows the sequence of the 1650 bp *Cla*I fragment, which contains 432 bases 5' to the beginning of the *pmi* coding region,
30 391 codons that contain the coding capacity for the *pmi* gene product (a protein of predicted Mr of 42.6 kDa), and 43 bases 3' to the TAG stop codon.

SEQ ID NOs:3 and 4 are two oligonucleotides that were prepared to facilitate cloning the *pmi* gene
35 from pADE253 into a plant transformation vector. The oligonucleotide of SEQ ID NO:3 is 20 bases in length and is identical to bases 391-410 of the *Cla*I fragment

(coding strand), except for position 403 where there is single mismatch. The oligonucleotide of SEQ ID NO:4 is 25 bases in length and is of complimentary sequence to bases 1641-1617 of the *Cla*I fragment, except for mismatches at positions 1628 and 1630.

The two oligonucleotides and DNA from plasmid pADE253 were used in a polymerase chain reaction (PCR) to amplify a population of approximately 1250 bp DNA fragments containing the *pmi* coding region and small portions of the upstream and downstream regions. As a result of the mismatched bases between the oligonucleotides and the *Cla*I fragment and because most of the DNA synthesized in the PCR reaction used newly synthesized DNA as a template rather than the pADE253 DNA as a template, the majority of the approximately 1250 bp fragments synthesized contained two new restriction endonuclease recognition sites that facilitated the cloning of the *pmi* gene into a plant transformation vector.

The new restriction endonuclease recognition sites added were a *Bgl*II recognition site (AGATCT), located between 24 and 29 bases 5' to the initiator methionine codon of the *pmi* gene, and a *Sac*I recognition site (GAGCTC), located between 21 and 26 bases 3' to the TAG stop codon of the *pmi* gene. The DNA from the PCR reaction was digested with *Bgl*II and *Sac*I for several hours at 37°C, and then electrophoresed on a 1 percent low melting point agarose gel. An approximately 1225 bp band was excised from the gel, and the DNA was isolated and purified.

This fragment, containing the *pmi* gene, was cloned into the plasmid pGEM3Zf(-) (Fig. 2). Plasmid pGEM3Zf(-) is a commercially available vector (Promega Corporation), and is 3199 bps in length. More specifically, pGEM3Zf(-) DNA was digested with *Bam*HI and *Sac*I, and the vector fragment was gel-purified. Digestion with *Sac*I leaves a four base, 3' overhang of

the sequence AGCT. Although the recognition sites for *Bam*HI and *Bgl*II are different, both enzymes produce an identical four base, 5' overhang of the sequence GATC. These overhanging regions greatly facilitate directional cloning.

The approximately 1.2 kilobase pair kbp *Bgl*II-*Sac*I fragment containing the *pmi* gene and the *Bam*HI and *Sac*I digested pGEM3Zf(-) DNA were ligated together overnight (about 18 hours) at 16°C. The following day, the ligation reaction product was used to transform *E. coli* strain MV1190. Various aliquots from the transformation were spread onto LB agar plates supplemented with carbenicillin (100 mg/L), isopropyl-1-thio- β -D-galactoside (IPTG) (0.1 mM), and 5-bromo-4-chloro-3-indolyl- β -D-galactoside (X-gal) (1 ml/L of a 20 mg/mL stock dissolved in N,N dimethylformamide).

Plasmid pGEM3Zf(-), like many commonly used plasmids in molecular biology, has a color marker selection system to indicate when DNA has been inserted into the vector's polylinker. When *E. coli* containing the omega fragment of β -galactosidase, which is usually contained on the F' plasmid, and harboring the pGEM3Zf(-) plasmid are plated onto medium supplemented with IPTG and X-gal, the α -fragment of β -galactosidase, which is encoded by the plasmid, can complement the omega fragment and produce a functional β -galactosidase enzyme. β -galactosidase can cleave the X-gal in the medium, yielding a blue color. Normally, when DNA is inserted into the polylinker region of pGEM3Zf(-), α -complementation is destroyed, and the colonies appear white instead of blue. The day following transformation, plasmid DNA was purified from several of the white colonies that grew on the LB-carbenicillin-IPTG-X-gal plates. To confirm the presence of the *pmi* gene, these plasmid DNAs were examined by their gel electrophoresis profiles after digestion with various restriction endonuclease

enzymes. A clone that contained the *pmi* gene was selected from this population and the plasmid was designated as pETO147 (Fig. 2).

5 The *pmi* gene was excised from plasmid pETO147 and inserted into a binary vector that facilitated transformation. pETO107 is approximately 13 kbp in length, and is a derivative of the commonly used binary plant transformation vector BIN19 [Bevan, Nucleic Acids Res., 12: 8711-8721(1984)]. Like BIN19, plasmid 10 pETO107 has both a right and left TDNA border region. These 25 base pair TDNA regions specify the sites that are recognized by *Agrobacterium tumefaciens*, which mediates the transfer of DNA from the plasmid to the plant chromosome; DNA between these borders is inserted 15 into the plant chromosome by *A. tumefaciens*. Infecting a number of different plant tissue types, such as leaf discs or cotyledons with *A. tumefaciens* that harbor a binary vector is a preferred method of transforming many plant species. Outside the TDNA borders, plasmid 20 pETO107 contains the bacterial *nptIII* gene, which can confer kanamycin resistance to *E. coli* that harbor this plasmid.

Plasmid pETO107 has two expression cassettes of interest between the TDNA borders. Starting from the 25 right TDNA border and proceeding clockwise for convenience (Fig. 2), the first cassette is composed of three elements that are operationally linked. The first element is the nopaline synthase promoter (P-Nos), which when inserted into the plant chromosome 30 directs the constitutive expression in most plant cells of genes that are positioned downstream of this DNA (clockwise in Fig. 2). The next element is the coding region for the neomycin phosphotransferase type II gene (NPTII). If NPTII is expressed in most plant cells in 35 sufficient quantities, it confers resistance to those cells to the antibiotic kanamycin. The third and final element of the first cassette is the 3' termination end

of the nopaline synthase coding region (Nos-T), which contains the polyadenylation recognition site. The first cassette is commonly used in plant transformation vectors as a selectable marker to identify transformed tissue; the kanamycin resistance conferred to transformed cells results in a selective growth advantage on medium supplemented with kanamycin. The *nptII* expression cassette is part of the original plasmid, and is not essential for the current invention, but is rather an example of a before-discussed second expression cassette.

Proceeding clockwise in Fig. 2, the second expression cassette in plasmid pETO107 also contains three elements. The first element is the 35S promoter (35S) from cauliflower mosaic virus (CaMV 35S). Like the P-Nos promoter, CaMV 35S directs constitutive expression in plant cells. Generally, the level of expression in plant cells is greater with CaMV 35S than P-Nos. The next element is a polylinker and includes the recognition sites for the following restriction endonuclease enzymes: *XbaI*, *BamHI*, *SmaI*, *KpnI* and *SacI*. The third and final element of this cassette is the 3' termination end of the nopaline synthase coding region (Nos-T). The polylinker region of this second element was used to insert the *XbaI*-*SacI* fragment from pETO147, which contains the *pmi* gene, into the *XbaI* and *SacI* sites that are located between the CaMV 35S and Nos-T elements.

DNA from plasmids pETO147 and pETO107 were digested with *XbaI* and *SacI*. The approximately 1.2 kbp *XbaI*-*SacI* fragment from plasmid pETO147, and the 13 kbp *XbaI*-*SacI* fragment from plasmid pETO107 were gel-purified. In a ligation reaction containing the 1.2 kbp *XbaI*-*SacI* fragment from plasmid pETO147 containing the *pmi* gene, and the 13 kbp *XbaI* and *SacI* plasmid pETO107 fragment, the 4-base overhangs facilitated the directional insertion of the *pmi* gene between the CaMV

35S promoter and Nos-T. The ligation reaction was performed overnight (about 18 hours) at 16°C.

The following day, the ligation reaction product was used to transform the *E. coli* strain XL1-Blue MRF'. Various aliquots from the transformation were spread onto LB agar plates supplemented with kanamycin (50 mg/L). The following day, plasmid DNA was purified from several colonies, and gel electrophoresis profiles from restriction endonuclease digestions were used to identify a plasmid containing a *pmi* gene that was inserted between the CaMV 35S and Nos-T elements. One such plasmid was designated pET0148 (Fig. 2).

Plasmid pET0148 was deposited at the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD on April 6, 1995 and was given ATCC accession number 97107. The present deposit was made in compliance with the Budapest Treaty requirements that the duration of the deposit should be for 30 years from the date of deposit or for five years after the last request for the deposit at the depository or for the enforceable life of a U.S. patent that matures from this application, whichever is longer. The vector will be replenished should it become non-viable at the depository.

E. coli harboring plasmid pET0148 were mobilized into the disarmed *A. tumefaciens* strain LBA4404 using the triparental mating system to form *A. tumefaciens* pET0148::LBA4404. Transconjugants were used to transform tomato (*Lycopersicon esculentum*).

To examine the dynamics of transformation and regeneration on mannose in several crops, a scorable marker was added to the plasmid pET0148 transformation vector. Specifically, an additional expression cassette, containing the CaMV 35S promoter, β -glucuronidase coding region and Nos-T was inserted into plasmid pET0148. The resultant vector, designated

plasmid pETO156, was mated into *A. tumefaciens* by the triparental method.

Example 1: Transformation of Tomato Plants

5 Tomato seeds were sterilized in a solution containing 20 percent of a commercial household bleach containing 5.25 percent sodium hypochlorite, rinsed three times in sterile distilled water and planted on Murashige and Skoog (MS) medium (Gibco) solidified with
10 10 grams of Noble agar (Gibco) in 135 mm Phytacorn™ tissue culture vessels (Sigma, St. Louis, MO). Seeds were germinated for 72 hours at 25°C in the dark, then moved to a lighted shelf under approximately 80 $\mu\text{mol m}^{-2}\text{s}^{-1}$ PPF, at 24-26°C. Plant tissue used for
15 transformation was prepared by removing cotyledons from 7-day-old seedlings and cutting them into three parts (proximal, middle and distal to the growing point).

The middle and proximal parts were used for co-cultivation with *Agrobacterium*. Those parts were
20 placed abaxial side down on a sterile filter paper overlaying co-cultivation medium R1F supplemented with 0.09 M (16 g/L) glucose, and incubated in the dark for 24 hours. Next, they were incubated for 20 minutes with bacterial inoculum containing 5×10^8 CFU/mL of
25 *Agrobacterium tumefaciens*, pETO148::LBA4404, blotted dry, and cultured for 48 hours, at 24°C, in the dark.

Bacterial inoculum was prepared by growing *A. tumefaciens*, pETO148::LBA4404, in 25 ml of AB medium [Chilton et al., 1974, Proc. Natl. Acad. Sci. USA,
30 71:3672-3676 (1974)] supplemented with 50 mg/L kanamycin (K) and 25 mg/L streptomycin (St) (AB_{K50St25}) on a shaker at 28°C, 180 rpm, for 24 hours. One mL of this culture was transferred to 25 mL of fresh AB_{K50St25} medium and grown on a shaker at 28°C, 180 rpm for 24 hours.
35 The bacteria were then pelleted by centrifuging at 8000 rpm for 10 minutes in a Beckman J2-21 centrifuge using a JA-20 rotor. The bacterial pellet was resuspended in

sterile MS medium and their concentration was adjusted to 5×10^8 CFU/mL using spectrophotometric optical density reading at 550 nm ($0.1 \text{ OD}_{550} = 2 \times 10^8$ CFU/mL). Prior to co-cultivation, the inoculum was supplemented with
5 acetosyringone (3',5'-dimethoxy-4'-hydroxy-acetophenone, Sigma, St. Louis, MO) to a final concentration of 600 micromolar.

Selection of Transformed Tomato Plants

10 After the 48 hour co-cultivation period with the *Agrobacterium* (pETO148::LBA4404) on the R1F medium, tomato tissue was moved to MR1 regeneration medium containing 0.09 M mannose as the sole carbon source. The MR1 medium was supplemented with 100 mg/L of the
15 antibiotic ticarcillin (Duchefa Co.) to eliminate *Agrobacterium*. After two weeks of culture, tomato tissue was moved to fresh MR1 medium. At this time small calli were visible on the cut ends of co-cultivated tissue, whereas control tissue that was not
20 co-cultivated with the bacteria but was also cultured on the MR1 medium, showed yellowing and did not produce any callus. After two more weeks of culture all tissues were moved to MR1/2 regeneration medium also containing 0.09 M mannose as a sole carbon source. At
25 this time the co-cultivated tissue produced a thick ridge of callus with small shoot primordia, whereas the control tissue was yellow in color, did not enlarge, nor produce callus. After two weeks on the MR1/2 medium the calli and shoot primordia that formed on the
30 co-cultivated tissue were detached from the original tissue and moved to MRO medium supplemented with 0.09M mannose. On this medium the shoots elongated and developed roots. Rooted plants were potted and moved to the greenhouse.

Biochemical and Molecular Analysis
of Regenerated Plants

Plants regenerated on mannose were assayed biochemically and molecularly to determine whether they were transformed. These plants were selected by their ability to grow on a medium containing mannose as an encrypted carbon source, but because the pET0148 transformation vector contained both the *pmi* and the *nptII* genes, plants were assayed for transformation indirectly by scoring for the expression of the second gene (*nptII*) using a commercially available (5 Prime → 3 Prime, Inc.) NPTII enzyme linked immunoabsorbant assay (ELISA).

Fig. 3 shows the relative NPTII expression from three rank-ordered populations. One population included five tomato plants that were non-transformed (negative control). The second population of five plants was from a transformation study that used kanamycin as a selective agent. In these positive control plants, the NPTII ELISA assays directly for the expression of the selective marker gene which detoxifies the selective agent. The expression of NPTII in these plants covered a wide range. The third population were 16 plants selected on mannose. The NPTII expression in the plants selected on mannose also covered a wide range. The expression of NPTII in the plants selected on mannose indicated that the *nptII* expression cassette that is incorporated with the *pmi* expression cassette was functioning in plants, and supplied indirect evidence that these plants were transformed. Further, these data indicate that the expression from the *pmi* gene, which confers the ability to grow on mannose, did not affect expression of an adjacent gene in the plant's chromosome.

To assay directly for transformation, a Southern blot was performed on several putative transformants. A Southern blot is a commonly performed assay in molecular biology that can show how many copies of the

transgene of interest are incorporated into the genome. Five grams of young leaf tissue were harvested and used to isolate genomic DNA. DNA yields were quantitated using a Hoefer Model TKO100 Fluorometer.

5 For each sample, 10 micrograms of DNA was digested with *HindIII*, electrophoresed on a 1 percent agarose gel and transferred to a nylon membrane. DNAs were fixed to the membrane using a Stratagene CL-100 ultraviolet crosslinker. The GeniusTM non-radioactive
10 detection system (Boehringer Mannheim) was used to probe the membrane and the blot results were recorded by fluorography. Using primers specific for the *pmi* gene, a 788 bp fragment was amplified using PCR. During amplification, digoxigenin-11-dUTP was
15 incorporated into the PCR fragments. The *pmi*-specific, digoxigen-labeled fragments were used to probe the genomic blot. Protocols for the hybridization, blot washing and visualization were supplied by the manufacturer.

20 A blot of 11 putative transgenic samples was prepared using molecular mass markers from *HindIII*-digested bacteriophage lambda DNA and DNA from a non-transformed tomato plant as a negative control. The fluorogram showed that the *pmi* probe did not hybridize
25 to the DNA from the non-transformed parental control, but did decorate specific bands in nine of the eleven putative transformants. The number of copies inserted into the tomato genome varied (data not shown).

30 Example 2: Monocot Cell Growth on Mannose

To assay whether mannose could be used for selection of transgenic cells in monocot plants the growth of cell suspensions of both oats and corn (maize) was evaluated in sucrose-supplemented media
35 (standard protocol) and in media in which sucrose was replaced with mannose at the same molar concentration.

Oats

Approximately 500 mg of drained oat suspension cells were inoculated to 40 ml of liquid MS2-D medium (4.4 g/L MS salts, Sigma #M5524; 0.5 mg/L thiamine HCl; 2 mg/L 2,4-D; 150 mg/L L-asparagine; pH=5.8) containing either 0.06 M sucrose (20 g/L) or 0.06 M mannose (10.6 g/L). Each treatment was replicated three times. The flasks were placed on a shaker at 160 rpm, 24-26°C, using 16 hours of photoperiod. After 25 days, the suspension cells were allowed to settle at the bottom of a test tube, the liquid medium was aspirated and the cells were weighed. The net weights of the cells are listed in Table 2, below.

15

Table 2

Net weight (grams) of Oat Suspension Cells After Culture in Media Supplemented Either with Sucrose or Mannose as Carbon Source

20

	<u>Replicate</u>	<u>Sucrose</u>	<u>Mannose</u>
	1	3.03	0.79
25	2	3.33	0.66
	3	3.41	0.92
30	Mean \pm SE	3.26 \pm 0.12	0.79 \pm 0.08

Corn (Maize)

A rapidly growing suspension culture of black Mexican sweet corn (BMS) was washed several times in mannose-supplemented (10.6 g/L) MS-F medium (4.4 g/L MS salts, Sigma #M5524; 1.3 mg/L nicotinic acid; 0.25 mg/L pyridoxine HCl; 0.25 mg/L thiamine HCl; 0.25 mg/L calcium pantothenate; 2 mg/L 2,4-D; 0.1 g/L inositol; 150 mg/L L-asparagine; pH=5.8). A five mL aliquots of a thick (1:1 cells to liquid ratio) cell slurry made from the washed cells were inoculated to

flasks containing 35 mL of either mannose-supplemented (0.06 M) or sucrose-supplemented (0.06 M) medium. Each treatment was replicated four times. The flasks were placed on a shaker at 160 rpm, 24-26°C, in the dark. The growth of the suspension cells was assessed after 32 days of culture by measuring the packed cell volume. The results are presented in Table 3, below.

Table 3

Volume (cc) of Corn Suspension Cells After Culture in Media Supplemented Either with Sucrose or Mannose

<u>Replicate</u>	<u>Sucrose</u>	<u>Mannose</u>
1	8.4	1.8
2	7.6	2.0
3	8.0	2.4
4	10.4	2.0
<hr/>		
Mean \pm SE	8.60 \pm 0.62	2.05 \pm 0.13

Both oat and corn cell suspensions showed severe inhibition of growth when cultured in mannose-supplemented medium as compared to the standard sucrose-containing medium. This indicates that a replacement of sucrose with mannose provides an effective selection for genetically engineered cells from monocot plants that can utilize mannose as a carbon source under heterotrophic growth conditions.

Example 3: Regeneration of transgenic tomatoes

Tomato tissue was prepared and cocultivated with *Agrobacterium* containing the *pmi* gene as described in Example 1. After cocultivation tomato tissue was cultured either on the standard MR1 medium

-58-

or on the MR1 medium supplemented with 170.00 mg/L of monobasic potassium phosphate (MR1/P medium). The supplementation of the mannose-containing medium with additional phosphorus resulted in a significant increase in the number of transformed plants that were regenerated. The comparison of regeneration on MR1 medium and MR1/P medium is shown in Table 4.

Table 4

Comparison of Regeneration of Transformed Plants on MR1 and MR1/P Media

	<u>Medium</u>	<u>Frequency of regeneration (Percent)</u>	<u>Number of regenerated shoots</u>	<u>Percent normal shoots</u>	<u>Percent abnormal shoots</u>
	MR1	66.4	83	48.2	51.8
	MR1/P	82.4	103	80.6	19.4

These observations indicate that an additional modification of the basic medium composition results in increased regeneration of transformed plants. Doubling of the concentration of phosphorus, which is an essential component of the regeneration medium, gave further growth advantage to tomato cells containing the *pmi* gene, resulting in a more efficient regeneration of genetically engineered plants using mannose for selection.

Example 4: Transgenic Melon

Decoated melon seeds were surface sterilized in a solution containing 10 percent of a commercial household bleach containing 5.25 percent sodium hypochlorite for ten minutes, rinsed six times in sterile, distilled water and germinated on a blotter paper soaked with sterile water for 24 hours, in the

dark. After germination, cotyledons were detached from the seed axis and cultured on B0 medium for two days under $20 \mu\text{mol m}^{-2}\text{s}^{-1}$ PPFD. Tissue for transformation was cut out using a No. 1 corkborer and inoculated with *Agrobacterium tumefaciens* pET0156::EHA105 by first briefly vortexing and then by soaking the tissue in the bacterial inoculum for 20 minutes. Inoculated tissue was cultured for three days on a fresh B0 medium overlaid with a sterile filter paper. Cocultivated explants were washed three times in a liquid B0 medium containing 1000 mg/L of the antibiotic carbenicillin and 200 mg/L of the antibiotic ticarcillin (Duchefa Co.) to eliminate the *Agrobacterium*, followed by two weeks of culture on shoot induction medium B300 supplemented with 16 gm/L of mannose, at 24-26°C, under 60-80 $\mu\text{mol m}^{-2}\text{s}^{-1}$ PPFD of light, at 16 hours of photoperiod.

After two weeks on the B300 medium melon tissue was transferred to shoot elongation medium E500 also containing 16 gm/L mannose, and cultured for another two weeks. At the time of transfer to the E500 medium the cocultivated tissue showed rapid cell proliferation, whereas non-transformed control tissue did not enlarge and did not produce any callus. Shoots that developed on the cocultivated tissue were rooted on medium N500 containing 16 gm/L mannose. Noncocultivated control tissue never produced any shoots. Rooted plants were potted. Effective selection of transformed melon plants was confirmed by staining plant tissue in the chemical reagent X-Gluc, which causes the transgenic tissue to turn blue due to the production of indigo dye by transgenic cells.

All media used in melon regeneration contained 4.3 gm/L of MS salts (Gibco #11117-074) and 1 ml/L of MS vitamins (Sigma #7150), were solidified with 8.0 gm/L of Noble agar and had pH adjusted to 5.8. Media B300, E500 and N500 contained 1000 mg/L carbenicillin

-60-

and 200 mg/L ticarcillin (Duchefa Co.) to eliminate the *Agrobacterium*. The B0 medium used for preculture and cocultivation was supplemented with 30 gm/L of sucrose, whereas media B300, E500 and N500 contained 16 gm/L mannose as a sole carbon source. Medium B300 contained 1.0 mg/l benzyladenine; medium E500 contained 0.15 mg/L benzyladenine and 3 mg/L gibberellic acid; medium N500 contained 50 mg/L 1-naphthylacetic acid and 16 mg/L thiamine HCl.

A preferred way to regenerate transgenic melon plants using mannose as a selective agent is to use B300, E500 and N500 media supplemented with additional 170 mg/L or 340 mg/L monobasic potassium phosphate.

Example 5: Transgenic Squash

Decoated squash seeds were surface sterilized in a solution containing ten percent of a commercial household bleach containing 5.25 percent sodium hypochlorite for ten minutes, rinsed five times in sterile, distilled water and germinated for 24 hours in the dark at 27°C on filter paper soaked with sterile water. Plant tissue used for transformation was prepared by removing the shoot axis cotyledonary node tissue and the distal half of the cotyledonary tissue. The remaining proximal half was cut into 4 explant pieces.

All explants were inoculated for 10 minutes in 2.5×10^8 CFU/mL of *Agrobacterium tumefaciens*, pE10156:EHA 105 in liquid MS medium supplemented with 0.2 mM acetosyringone. Explants were blotted on sterile filter paper and cocultivated adaxial side down on MS medium supplemented with 2 mg/L 2,4,5-T and 0.5 mg/L kinetin. Explants were cocultivated for 3 days in the dark in unsealed Petri dishes.

After the 3 day cocultivation period, explants were washed for 3 hours in MS liquid medium supplemented 10 mg/L 2,4-D, 0.5 mg/L kinetin, 400

mg/l cefotaxime and 750 mg/L carbenicillin. The
rinse medium was changed 3 times during the
procedure. Explants were blotted on sterile filter
paper and cultured abaxial side down on selection
5 medium consisting of MS salts and vitamins, 10 mg/L
2,4-D, 0.5 mg/L kinetin, 16.2 g/L mannose, 1000 mg/L
carbenicillin and 170 mg/L monobasic potassium
phosphate. Explants were cultured in unsealed Petri
dishes and kept in diffused light. Explants were
10 subcultured to fresh medium every two weeks.

After 8 weeks of selection, clusters of
embryos formed on the edges of the explants cultured
on mannose, whereas control explants did not produce
any embryos. The transgenic character of the
15 regenerating embryos was confirmed by staining in the
chemical reagent X-Gluc, which causes transgenic
tissue to turn blue. Regeneration of transgenic
plants from embryos was achieved by subculture to
hormone-free medium for rooting and further
20 development. The transgenic character of the plants
regenerated on mannose-containing medium was
confirmed by staining in Xgluc.

Example 6: Transgenic Squash

Two experiments were performed in squash
25 using the mannose-based selection. In both
experiments the preparation of the explants and
their inoculation was as follows:

Decoated squash seeds were surface
sterilized in a solution containing twenty percent
30 of a commercial household bleach containing 5.25
percent sodium hypochlorite (Clorox™) for ten
minutes, rinsed briefly five times in sterile
distilled water and germinated for 24 hours in the
dark at 27°C on filter paper soaked with sterile
35 water. Plant tissue used for transformation was
prepared by removing the shoot axis, cotyledonary
node tissue and the distal half of the cotyledonary

-62-

tissue. The remaining proximal half was cut into 4 pieces, which are referred to as explants.

The explants were inoculated for 10 minutes in 10 ml per 32 explants of MS medium supplemented with 0.2 mM acetosyringone, and 2.5×10^8 CFU/ml of the disarmed *Agrobacterium tumefaciens*, strain EHA 105, that was harboring the binary plasmid pETO156. After incubation with the *Agrobacterium*, the explants were blotted on sterile filter paper and placed adaxial side down on MS medium supplemented with 2 mg/l 2,4,5-T and 0.5 mg/l kinetin. Explants were cultured for 3 days in unsealed Petri dishes, in the dark.

After the 3 days of culture, explants were washed for 3 hours in 20-30 ml of MS liquid medium supplemented with 10 mg/l 2,4-D, 0.5 mg/l kinetin, 400 mg/l cefotaxime and 750 mg/l carbenicillin. The rinse medium was changed every hour for a total of 3 washes. After washing, explants were blotted on sterile filter paper and placed abaxial side down in Petri plates containing selection medium consisting of MS salts and vitamins, 10 mg/l 2,4-D, 0.5 mg/l kinetin, 1000 mg/l carbenicillin.

In the first experiment the medium contained 16.2 g/l mannose and various concentrations of KH_2PO_4 (170 mg/l, 340 mg/l or 510 mg/l). Explants were cultured in unsealed Petri dishes and kept in diffused light of approximately $10 \mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ Photosynthetic Photon Flux Density (PPFD). Explants were subcultured to fresh medium every two weeks.

In the second experiment explants were cultured on the MS medium supplemented with the following:

- 16.2 g/l mannose,
- 16.2 g/l mannose plus 170 mg/l KH_2PO_4 ,

-63-

30 g/l sucrose for one week, then selection on
 16.2 g/l mannose plus 170 mg/l KH_2PO_4 ,
 30 g/l sucrose for two weeks, then selection on
 16.2 g/l mannose plus 170 mg/l KH_2PO_4 .

5 In both experiments clusters of embryos
 formed on the edges of the cocultivated with
 Agrobacterium explants after 8 weeks of culture,
 whereas non-cocultivated control explants did not
 produce embryos. Transformation of the regenerating
 10 embryos was confirmed by assaying for the expression
 of the GUS gene, which causes transgenic tissue to
 turn blue in the presence of the chemical 5-Bromo-4-
 Chloro-3-Indolyl- β -D-Glucuronic Acid
 Cyclohexylammonium Salt (X-Gluc). Because the GUS
 15 gene contains an intron, expression of this gene can
 occur in the eukaryotic cells, but not in the cells
 of the prokaryotic Agrobacterium. Explants were
 scored as transgenic if a blue signal was observed
 in the embryos. The percentage of explants
 20 producing transgenic embryos in the first experiment
 is shown in Table 5 and the percentage of explants
 producing transgenic embryos in the second
 experiment is shown in Table 6.

25 Table 5. Effect of Phosphate Concentration on the
 Selection of Squash Embryos Transformed with the
pmi Gene.

Treatment	% Explants with Transformed Embryos
MS medium + 16.2 g/l mannose	2.8
MS medium + 16.2 g/l mannose + 170 mg/l KH_2PO_4	13.9
MS medium + 16.2 g/l mannose + 340 mg/l KH_2PO_4	16.7
MS medium + 16.2 g/l mannose + 510 mg/l KH_2PO_4	5.6

Table 6. Effect of Carbon and Phosphate on the Selection of Squash Embryos Transformed with the *pmi* Gene.

5	Treatment	% Explants with Transformed Embryos
	MS medium + 16.2 g/l mannose	4.3
	MS medium + 16.2 g/l mannose + 170 mg/l KH_2PO_4	8.3
10	MS medium + 30 g/l sucrose for 1 week, then 16.2 g/l mannose + 170 mg/l KH_2PO_4	8.5
15	MS medium + 30 g/l sucrose for 2 weeks, then 16.2 g/l mannose + 170 mg/l KH_2PO_4	15.5

In the first experiment the frequency of transgenic embryo production using mannose selection was increased by 5-fold if the concentration of phosphate in the selection medium was doubled and 6-fold if the concentration was tripled when compared to the control (MS medium) concentration of phosphate. The MS medium is a widely used formulation in the regeneration of many plants from tissue culture.

In the second experiment the use of mannose plus phosphate for selection resulted in a nearly two-fold increase in transformation frequency. When the explants were first grown on an MS medium containing 30 g/l sucrose and transferred a week later to an MS medium containing 16.2 g/l mannose and 170.00 mg/l KH_2PO_4 , no significant increase in the transformation frequency was observed over the explants kept on MS + mannose + KH_2PO_4 medium. In contrast, a two week growth on MS medium supplemented with sucrose followed by growth on MS + mannose + KH_2PO_4 medium resulted in a nearly two-fold increase in transformation frequency over explants grown on MS + mannose + KH_2PO_4 medium.

-65-

Further, the two week sucrose treatment followed by the mannose + KH_2PO_4 medium displayed a 3.5-fold increase in transformation efficiency over the mannose only treatment.

5 Both experiments showed that the addition of phosphate to the MS-mannose medium increased the frequency of regeneration of transgenic embryos from 3.5-fold up to 6-fold. The second experiment also showed that delay in the selection by up to two
10 weeks resulted in an increased recovery of transgenic embryos. The differences in the frequencies of transformation in these two experiments were characteristic of a high experiment to experiment variation frequently observed in
15 tissue culture.

Example 7. Transgenic Tomato.

Seeds from two transgenic tomato lines, X11-28 and X11-40, transformed with the *pmi* gene
20 were sterilized in 20% Clorox™ for 20 minutes, rinsed 3 times in sterile distilled water and planted on Murashigi and Skoog medium (Gibco) solidified with 7 g/L of commercial grade Phytagar in 135 mm Phytacorn™ tissue culture vessels (Sigma,
25 St. Louis, MO). Seeds were germinated for 72 hours at 25°C in the dark, then moved to a lighted shelf under approximately 80 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD, at 24-26°C. Plant tissue used in the experiment was prepared by removing cotyledons from 7-day-old seedlings and
30 cutting them into three sections, proximal, middle and distal to the growing point. The middle and proximal parts were placed either on the control R1 medium supplemented with 16g/L glucose or on the MR1 regeneration medium containing 16 g/L mannose.
35 Both the R1 and MR1 media were subdivided into three treatments, which differed in the amount of monobasic potassium phosphate: 1x MS (standard concentration of 170 mg/L KH_2PO_4), 2x MS (340 mg/L

-66-

total phosphate concentration) and 3x MS (510 mg/L total phosphate concentration). Each treatment was replicated three times, each replicate consisted of 9 explants. Explants were cultured for three weeks under approximately 30 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD, at 24-26°C, 16 hr photoperiod, then moved to a lighted shelf under approximately 80 $\mu\text{mol}\cdot\text{m}^{-2}\cdot\text{s}^{-1}$ PPFD. The effect of the phosphate concentration and carbon source on the growth of the explants was assessed after 6 weeks of culture by weighing each explant. The results are presented in Tables 7 and 8.

Table 7. Fresh weight (mg) of tomato explants from transgenic tomato line X11-28 transformed with the *pmi* gene cultured on two types of carbon and three levels of phosphate.

KH ₂ PO ₄ level	Carbon source	Replicate 1	Replicate 2	Replicate 3	Mean \pm Standard Error	Percent Difference
20 1x (standard)	Glucose	276.6	335.4	450.3	354.1 \pm	
	Mannose	251.3	362.9	346.5	320.3 \pm	- 9.5
2x	Glucose	221.3	296.0	392.2	303.1 \pm	
	Mannose	433.9	391.8	380.8	402.2 \pm	+ 32.7
25 3x	Glucose	258.5	323.5	240.7	274.3 \pm	
	Mannose	370.4	340.3	420.0	376.9 \pm	+ 37.4

Table 8. Fresh weight (mg) of tomato explants from transgenic tomato line X11-40 transformed with the *pmi* gene cultured on two types of carbon and three levels of phosphate.

K ₂ PO ₄ level	Carbon source	Replicate 1	Replicate 2	Replicate 3	Mean \pm Standard Error	Percent Difference
1x (standard)	Glucose	312.7	515.1	404.8	410.9 \pm	
	Mannose	441.1	550.1	382.3	457.8 \pm	+ 11.1
2x	Glucose	265.2	307.3	332.9	301.8 \pm	
	Mannose	412.8	384.1	388.4	395.1 \pm	+ 30.9
3x	Glucose	271.8	390.2	nt	331.0 \pm	
	Mannose	457.7	404.9	nt	431.3 \pm	+ 30.3

Tomato explants transformed with the *pmi* gene had on average 30% higher fresh weight when cultured on the mannose medium supplemented with two- or three-fold higher levels of phosphate as compared to medium containing glucose as a carbon source. This result indicates that elevated phosphate provides a growth advantage for transgenic cells transformed with the mannose isomerase gene on mannose medium.

In a transformation experiment, there are many nontransgenic cells and a few transgenic cells. The purpose of selection is to identify and provide a growth advantage to the transgenic cells with totipotency to regenerate plants. Cells derived either from photosynthetically active tissues, such as leaves, or storage organs, such as cotyledons of germinating seeds, are typically rich in metabolizable carbon. In a transformation experiment with the mannose isomerase gene and the use of mannose medium, both the nontransgenic cells, which are present in great excess, and the transgenic cells, which are relatively scarce, have the potential to grow by using the residual intracellular supply of carbon. The transgenic

-68-

cells have an additional source of carbon available to them because they can access the carbon that is present in the medium in the form of mannose. If the amount of growth due to the residual intracellular supply of carbon is significant, it will be more difficult to identify the few cells that are transgenic.

Data shown in Tables 7 and 8 demonstrate that the addition of phosphate confers a growth advantage to transgenic cells grown on mannose over transgenic cells grown on glucose. Further, these experiments suggest why transformation experiments on mannose medium supplemented with elevated phosphate levels are far more successful than experiments with mannose only. In the mannose plus phosphate environment, the transgenic cells have two growth advantages over the glucose-utilizing (i.e. nontransgenic) cells: (1) they can access additional carbon from the medium, and (2) the phosphate confers an additional growth advantage to cells capable of growing on mannose over cells utilizing glucose. This double growth advantage results in an efficient selection of transgenic cells that have the totipotency to regenerate healthy transgenic plants.

The foregoing is intended as illustrative of the present invention but not limiting. Numerous variations and modifications can be effected without departing from the true spirit and scope of the novel concepts of the invention.

-69-

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
- 5 (i) APPLICANT: Trulson, Anna J.
Green, Charles E.
Braun III, Carl J.
- 10 (ii) TITLE OF INVENTION: PROCESS FOR SELECTION OF TRANSGENIC
PLANT CELLS
- (iii) NUMBER OF SEQUENCES: 4
- 15 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Greer, Burns & Crain, Ltd.
(B) STREET: 233 South Wacker Drive, Suite 8660
(C) CITY: Chicago
(D) STATE: Illinois
20 (E) COUNTRY: USA
(F) ZIP: 60606-4002
- (v) COMPUTER READABLE FORM:
25 (A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
30 (A) APPLICATION NUMBER: US 08/417,618
(B) FILING DATE: 06-APR-1995
(C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
35 (A) NAME: Mueller, Lisa V.
(B) REGISTRATION NUMBER: 38,978
(C) REFERENCE/DOCKET NUMBER: 1605.61148
- (ix) TELECOMMUNICATION INFORMATION:
40 (A) TELEPHONE: (312) 993-0080
(B) TELEFAX: (312) 993-0633
- (2) INFORMATION FOR SEQ ID NO:1:
- 45 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1650 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
50 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- 55 (ix) FEATURE:
(A) NAME/KEY: CDS
(B) LOCATION: 432..1607
- 60 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

- 70 -

ATCGATAACT TTCCACGCGA TGTCGCAGAG CTGGTGGACT GGTTCGACGC TCGCGACCCT
 60
 AACCGCATGT GCGCCCGGTG CCCGCTACGC GAGCAGATCC CGGTCTGGCT GTTGGGATCT
 5 120
 CTTGTCCTTC GAATTCGCGC ACGGAAACAT GTTCGCTGGT CAACAAGTAG TACTCGGTAT
 180
 10 CGTCCCTTTT GAGGGGAAAA GGGTCTTGAT AAAAGAAGGG TTTGTTGAC ATTGTGCTCT
 240
 CACTTACCGC TCGGTATGGT TATTCTCTGG GCAGGTGTTT CATTGCCCGA CTCAAAGCGA
 300
 15 GTAACACTAT CCTACACAAT TTTTAAACAA AAAGTACGAC AAGTACGACT TTTTACGCC
 360
 GGAGGTACT TCATCGGGT TTCTTGGTTT AATACCTCCC ATTGATCTCC ACATTGAAAC
 20 420
 AGGGCTTGAT A ATG CAA AAA CTC ATT AAC TCA GTG CAA AAC TAT GCC TGG
 470
 Met Gln Lys Leu Ile Asn Ser Val Gln Asn Tyr Ala Trp
 25 1 5 10
 GGA AGT AAA ACT GCG TTA ACG GAA CTT TAT GGC ATC GCC AAT CCG CAG
 518
 Gly Ser Lys Thr Ala Leu Thr Glu Leu Tyr Gly Ile Ala Asn Pro Gln
 30 15 20 25
 CAG CAG CCA ATG GCT GAA CTC TGG ATG GGC GCG CAT CCC AAA AGC AGC
 566
 Gln Gln Pro Met Ala Glu Leu Trp Met Gly Ala His Pro Lys Ser Ser
 35 30 35 40 45
 TCG CGA ATC ACC ACC GCC AAC GGC GAA ACC GTC TCC CTG CGT GAC GCC
 614
 Ser Arg Ile Thr Thr Ala Asn Gly Glu Thr Val Ser Leu Arg Asp Ala
 40 50 55 60
 ATC GAA AAG AAT AAA ACC GCC ATG CTG GGC GAA GCG GTA GCC AAC CGT
 662
 Ile Glu Lys Asn Lys Thr Ala Met Leu Gly Glu Ala Val Ala Asn Arg
 45 65 70 75
 TTC GGC GAA CTG CCG TTT CTG TTT AAA GTA CTG TGC GCC GCC AAA CCG
 710
 Phe Gly Glu Leu Pro Phe Leu Phe Lys Val Leu Cys Ala Ala Lys Pro
 50 80 85 90
 CTC TCT ATT CAG GTG CAC CCG AAT AAA CGC AAC TCC GAA ATC GGT TTC
 758
 Leu Ser Ile Gln Val His Pro Asn Lys Arg Asn Ser Glu Ile Gly Phe
 55 95 100 105
 GCG AAA GAA AAT GCG GCG GGT ATC CCC ATG GAT GCC GCA GAG CGG AAC
 806
 Ala Lys Glu Asn Ala Ala Gly Ile Pro Met Asp Ala Ala Glu Arg Asn
 60 110 115 120 125

-71-

TAT AAA GAT CCT AAC CAT AAA CCA GAG CTG GTT TTT GCC CTG ACG CCT
 854
 Tyr Lys Asp Pro Asn His Lys Pro Glu Leu Val Phe Ala Leu Thr Pro
 130 135 140

5
 TTC CTG GCG ATG AAC GCG TTC GCG GAA TTT TCT GAC ATT GTC TCT TTA
 902
 Phe Leu Ala Met Asn Ala Phe Arg Glu Phe Ser Asp Ile Val Ser Leu
 145 150 155

10
 CTG CAA CCT GTC GCC GGC GCG CAT TCC GCT ATC GCC CAC TTT TTG CAG
 950
 Leu Gln Pro Val Ala Gly Ala His Ser Ala Ile Ala His Phe Leu Gln
 160 165 170

15
 GTG CCG AAT GCT GAA CGT CTG AGC CAG CTT TTC GCC AGC CTG TTG AAT
 998
 Val Pro Asn Ala Glu Arg Leu Ser Gln Leu Phe Ala Ser Leu Leu Asn
 175 180 185

20
 ATG CAA GGC GAA GAA AAA TCC CGC GCG TTA GCC GTA CTC AAA GCG GCG
 1046
 Met Gln Gly Glu Glu Lys Ser Arg Ala Leu Ala Val Leu Lys Ala Ala
 190 195 200 205

25
 CTT AAC AGC CAG CAA GGC GAA CCG TGG CAA ACG ATC CGC GTG ATT TCA
 1094
 Leu Asn Ser Gln Gln Gly Glu Pro Trp Gln Thr Ile Arg Val Ile Ser
 210 215 220

30
 GAG TAT TAT CCT GAC GAC AGC GGG CTT TTC TCT CCT TTG TTG CTG AAT
 1142
 Glu Tyr Tyr Pro Asp Asp Ser Gly Leu Phe Ser Pro Leu Leu Leu Asn
 225 230 235

35
 GTG GTC AAA CTG AAT CCC GGC GAG GCG ATG TTC CTG TTT GCT GAA ACG
 1190
 Val Val Lys Leu Asn Pro Gly Glu Ala Met Phe Leu Phe Ala Glu Thr
 240 245 250

40
 CCT CAT GCT TAT CTG CAG GGC GTT GCG CTG GAA GTC ATG GCG AAC TCC
 1238
 Pro His Ala Tyr Leu Gln Gly Val Ala Leu Glu Val Met Ala Asn Ser
 255 260 265

45
 GAT AAC GTT CTG CGC GCT GGC CTT ACG CCA AAA TAT ATC GAC ATC CCT
 1286
 Asp Asn Val Leu Arg Ala Gly Leu Thr Pro Lys Tyr Ile Asp Ile Pro
 270 275 280 285

50
 GAG CTG GTC GCG AAC GTG AAG TTC GAA CCT AAG CCT GCC GGC GAG TTG
 1334
 Glu Leu Val Ala Asn Val Lys Phe Glu Pro Lys Pro Ala Gly Glu Leu
 290 295 300

55
 CTG ACT GCC CCG GTG AAA AGC GGC GCG GAG CTG GAC TTC CCA ATT CCG
 1382
 Leu Thr Ala Pro Val Lys Ser Gly Ala Glu Leu Asp Phe Pro Ile Pro
 305 310 315

- 72 -

GTT GAC GAT TTT GCT TTT TCA CTG CAC GAC CTG GCG CTT CAG GAG ACG
 1430
 Val Asp Asp Phe Ala Phe Ser Leu His Asp Leu Ala Leu Gln Glu Thr
 320 325 330

5 AGC ATC GGC CAA CAC AGC GCC GCG ATT CTG TTC TGC GTT GAG GGT GAG
 1478
 Ser Ile Gly Gln His Ser Ala Ala Ile Leu Phe Cys Val Glu Gly Glu
 335 340 345

10 GCG GTG TTA CGT AAA GAT GAA CAG CGT CTG GTA CTG AAG CCG GGT GAA
 1526
 Ala Val Leu Arg Lys Asp Glu Gln Arg Leu Val Leu Lys Pro Gly Glu
 350 355 360 365

15 TCT GCC TTT ATC GGC GCG GAT GAG TCT CCG GTT AAC GCC AGC GGC ACG
 1574
 Ser Ala Phe Ile Gly Ala Asp Glu Ser Pro Val Asn Ala Ser Gly Thr
 370 375 380

20 GGC CGT TTA GCG CGT GTT TAT AAC AAG CTG TAGCAACGTA CTGAATTTT
 1624
 Gly Arg Leu Ala Arg Val Tyr Asn Lys Leu
 385 390

25 TAACAACTCT TGCTAAGCTT ATCGAT
 1650

30 (2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 391 amino acids
 (B) TYPE: amino acid
 35 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

40 Met Gln Lys Leu Ile Asn Ser Val Gln Asn Tyr Ala Trp Gly Ser Lys
 1 5 10 15
 Thr Ala Leu Thr Glu Leu Tyr Gly Ile Ala Asn Pro Gln Gln Gln Pro
 45 20 25 30
 Met Ala Glu Leu Trp Met Gly Ala His Pro Lys Ser Ser Ser Arg Ile
 35 40 45
 50 Thr Thr Ala Asn Gly Glu Thr Val Ser Leu Arg Asp Ala Ile Glu Lys
 50 55 60
 Asn Lys Thr Ala Met Leu Gly Glu Ala Val Ala Asn Arg Phe Gly Glu
 65 70 75 80
 55 Leu Pro Phe Leu Phe Lys Val Leu Cys Ala Ala Lys Pro Leu Ser Ile
 85 90 95
 Gln Val His Pro Asn Lys Arg Asn Ser Glu Ile Gly Phe Ala Lys Glu
 60 100 105 110

- 73 -

Asn Ala Ala Gly Ile Pro Met Asp Ala Ala Glu Arg Asn Tyr Lys Asp
 115 120 125
 5 Pro Asn His Lys Pro Glu Leu Val Phe Ala Leu Thr Pro Phe Leu Ala
 130 135 140
 Met Asn Ala Phe Arg Glu Phe Ser Asp Ile Val Ser Leu Leu Gln Pro
 145 150 155 160
 10 Val Ala Gly Ala His Ser Ala Ile Ala His Phe Leu Gln Val Pro Asn
 165 170 175
 Ala Glu Arg Leu Ser Gln Leu Phe Ala Ser Leu Leu Asn Met Gln Gly
 180 185 190
 15 Glu Glu Lys Ser Arg Ala Leu Ala Val Leu Lys Ala Ala Leu Asn Ser
 195 200 205
 Gln Gln Gly Glu Pro Trp Gln Thr Ile Arg Val Ile Ser Glu Tyr Tyr
 20 210 215 220
 Pro Asp Asp Ser Gly Leu Phe Ser Pro Leu Leu Leu Asn Val Val Lys
 225 230 235 240
 25 Leu Asn Pro Gly Glu Ala Met Phe Leu Phe Ala Glu Thr Pro His Ala
 245 250 255
 Tyr Leu Gln Gly Val Ala Leu Glu Val Met Ala Asn Ser Asp Asn Val
 260 265 270
 30 Leu Arg Ala Gly Leu Thr Pro Lys Tyr Ile Asp Ile Pro Glu Leu Val
 275 280 285
 Ala Asn Val Lys Phe Glu Pro Lys Pro Ala Gly Glu Leu Leu Thr Ala
 35 290 295 300
 Pro Val Lys Ser Gly Ala Glu Leu Asp Phe Pro Ile Pro Val Asp Asp
 305 310 315 320
 40 Phe Ala Phe Ser Leu His Asp Leu Ala Leu Gln Glu Thr Ser Ile Gly
 325 330 335
 Gln His Ser Ala Ala Ile Leu Phe Cys Val Glu Gly Glu Ala Val Leu
 340 345 350
 45 Arg Lys Asp Glu Gln Arg Leu Val Leu Lys Pro Gly Glu Ser Ala Phe
 355 360 365
 Ile Gly Ala Asp Glu Ser Pro Val Asn Ala Ser Gly Thr Gly Arg Leu
 50 370 375 380
 Ala Arg Val Tyr Asn Lys Leu
 385 390

55

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

60

-74-

(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

10 AATACCTCCC ATAGATCTCC
20

(2) INFORMATION FOR SEQ ID NO:4:

15 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

20 (ii) MOLECULE TYPE: DNA (genomic)

25 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CTTAGCAAGA GCTCTTAAAA AATTC
25

30

We claim:

1. A process for selectively increasing the number of transformed plants regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions, the method comprising the steps of:
 - (a) culturing a mixture of transformed and non-transformed plant cells under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and growth by non-transformed plant cells except for a source of carbon that supports growth and proliferation and about 1.5 to 3 times the standard amount of phosphorus, said source of carbon being replaced by an encrypted carbon source that does not support growth and proliferation of said non-transformed cells, said transformed cells having a heterologous genomic DNA segment that contains at least one expression cassette,
 - the one expression cassette containing a heterologous DNA selectable marker segment that includes (i) a heterologous gene that encodes a heterologous enzyme that on expression converts said encrypted carbon source into a carbon source that supports growth and proliferation of said transformed plant cells under heterotrophic culture conditions, said first gene being operatively linked to (ii) a promoter DNA segment that controls expression of said heterologous gene, and (iii) a termination DNA segment;
 - (b) maintaining said heterotrophic culture conditions for a time period sufficient for said transformed plant cells to express said heterologous enzyme, grow and proliferate; and
 - (c) recovering said transgenic proliferating cells.

2. The process according to claim 1 wherein said heterologous gene is phosphomannose isomerase and said encrypted carbon source is mannose.

5

3. The process according to claim 1 wherein said phosphorous is a water-soluble phosphate salt.

4. The process of claim 3 wherein the phosphate salt is selected from the group consisting of: potassium phosphate, sodium phosphate, calcium phosphate, sodium hexametaphosphate, ammonium phosphate, orthophosphoric acid, ferric phosphate, sodium glycerophosphate, ferric glycerophosphate and organic phosphate.

5. The process according to claim 1 wherein expression from said promoter DNA sequence is repressed by a product of the normal metabolism of said transgenic plants under autotrophic growth.

6. The process according to claim 5 wherein the repressible promoter DNA sequence is that of the cucumber isocitrate lyase promoter.

25

7. The process according to claim 5 wherein the repressible promoter DNA sequence is that of the rice α -amylase Amy3A promoter.

8. The process according to claim 1 wherein the promoter is a constitutive promoter.

9. The process according to claim 8 wherein the constitutive promoter is selected from the group consisting of: the 35S promoter of the cauliflower

35

mosaic virus, the octopine synthase promoter and the nopaline synthase promoter.

10. A process for selectively increasing the
5 number of transformed plants regenerated from a mixture of transformed and non-transformed plant cells cultured under heterotrophic culture conditions, the method comprising the steps of

(a) culturing a mixture of transformed and
10 non-transformed plant cells under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and growth by non-transformed plant cells except for a source of carbon that supports growth and
15 proliferation and about 1.5 to 3 times the standard amount of phosphorus, said source of carbon being replaced by an encrypted carbon source that does not support growth and proliferation of said non-transformed cells, said transformed cells containing
20 a heterologous genomic DNA segment that contains at least one expression cassette,

the one expression cassette containing a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes
25 a heterologous enzyme that on expression converts said encrypted carbon source into a carbon source that supports growth and proliferation of said transformed plant cells under heterotrophic culture conditions, said first gene being operatively linked
30 to (ii) a first promoter DNA segment that controls expression of said heterologous gene, and (iii) a termination DNA segment;

(b) maintaining said heterotrophic culture conditions for a time period sufficient for said
35 transformed plant cells to express said heterologous enzyme, grow and proliferate;

-78-

- (c) recovering said proliferating cells; and
- (d) regenerating plant meristematic tissues or plant embryos from said proliferating cells.

5 11. The process according to claim 10 wherein said heterologous gene is phosphomannose isomerase and said encrypted carbon source is mannose.

10 12. The process according to claim 10 wherein said phosphorous is a water-soluble phosphate salt.

15 13. The process of claim 12 wherein the phosphate salt is selected from the group consisting of: potassium phosphate, sodium phosphate, calcium phosphate, sodium hexametaphosphate, ammonium phosphate, orthophosphoric acid, ferric phosphate, sodium glycerophosphate, ferric glycerophosphate and organic phosphate.

20 14. The process according to claim 10 wherein expression from said promoter DNA sequence is repressed by a product of the normal metabolism of said transgenic plants under autotrophic growth.

25 15. The process according to claim 14 wherein the repressible promoter DNA sequence is that of the cucumber isocitrate lyase promoter.

30 16. The process according to claim 14 wherein the repressible promoter DNA sequence is that of the rice α -amylase Amy3A promoter.

35 17. The process according to claim 10 wherein the promoter is a constitutive promoter.

-79-

18. The process according to claim 17 wherein the constitutive promoter is selected from the group consisting of: the 35S promoter of the cauliflower mosaic virus, the octopine synthase promoter and the nopaline synthase promoter.

19. A process for selectively increasing the number of transformed plants regenerated from a mixture of transformed and non-transformed plant cells placed under selective heterotrophic culture conditions, the method comprising the steps of:

(a) culturing a mixture of transformed and non-transformed plant cells for up to two weeks in a first culture medium that contains the minimal nutrients required for proliferation and growth by non-transformed plant cells including a source of carbon that supports growth and proliferation, said transformed plants cells containing a genomic heterologous DNA segment that contains at least one expression cassette,

the one expression cassette containing a heterologous DNA selectable marker segment that includes (i) a heterologous gene that encodes a heterologous enzyme that on expression converts an encrypted carbon source into a carbon source that supports growth and proliferation of said transformed plant cells under heterotrophic culture conditions, said first gene being operatively linked to (ii) a promoter DNA segment that controls expression of said heterologous gene, and (iii) a termination DNA segment;

(b) removing the mixture of transformed and non-transformed plant cells from the first culture medium;

35

-80-

- (c) placing the transformed and non-transformed plant cells under heterotrophic culture conditions in a second culture medium that contains the minimal nutrients required for proliferation and growth of the non-transformed plant cells except for an encrypted carbon source that does not support growth and proliferation of said non-transformed plant cells and 1.5 to 3 times the standard amount of phosphorous;
- (d) maintaining said heterotrophic culture conditions for a time period sufficient for said transformed plant cells to express said heterologous enzyme, grow and proliferate; and
- (e) recovering said proliferating cells.

15

20. The process according to claim 19 wherein said heterologous gene is phosphomannose isomerase and said encrypted carbon source is mannose.

20

21. The process according to claim 19 wherein said phosphorous is a water-soluble phosphate salt.

25

22. The process according to claim 21 wherein the phosphate salt is selected from the group consisting of: potassium phosphate, sodium phosphate, calcium phosphate, sodium hexametaphosphate, ammonium phosphate, orthophosphoric acid, ferric phosphate, sodium glycerophosphate, ferric glycerophosphate and organic phosphate.

30

23. The process according to claim 19 wherein expression from said promoter DNA sequence is repressed by a product of the normal metabolism of said transgenic plants under autotrophic growth.

35

-81-

24. The process according to claim 23 wherein the repressible promoter DNA sequence is that of the cucumber isocitrate lyase promoter.

5 25. The process according to claim 23 wherein the repressible promoter DNA sequence is that of the rice α -amylase Amy3A promoter.

10 26. The process according to claim 19 wherein the promoter is a constitutive promoter.

 27. The process according to claim 26 wherein the constitutive promoter is selected from the group consisting of: the 35S promoter of the cauliflower mosaic virus, the octopine synthase promoter and the nopaline synthase promoter.

15

 28. A process for selectively growing twice-transformed plants from a mixture of twice- and once-transformed plant cells comprising the steps of:

20

(a) culturing a mixture of twice- and once-transformed plant cells under heterotrophic culture conditions in a culture medium that contains minimal nutrients required for proliferation and growth by said once-transformed plant cells except for a source of carbon that supports growth and proliferation of said once-transformed cells and about 1.5 to 3 times the standard amount of phosphorus, said source of carbon being replaced by a second encrypted carbon source that does not support growth and proliferation of said once-transformed plant cells; said twice-transformed cells containing first and a second heterologous DNA segments that contains at least two expression cassettes, wherein at least one expression cassette

25

30

35

is in the first heterologous DNA segment and at least one expression cassette is in the second heterologous DNA segment;

the expression cassette in the first hetero-
5 logous DNA segment containing a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts a first encrypted carbon source into a carbon source that supports growth and
10 proliferation of said once- and twice-transformed plant cells under heterotrophic culture conditions but does not support growth and proliferation of non-transformed plant cells, said first gene being operatively linked to (ii) a first promoter DNA
15 segment that controls expression of said heterologous gene, and (iii) a termination DNA segment;

the expression cassette in the second hetero-
logous DNA segment containing a second heterologous
20 DNA selectable marker segment that includes (i) a second heterologous gene that encodes a second heterologous enzyme that on expression during heterotrophic culture of said twice-transformed cells converts said second encrypted carbon source
25 that does not support growth and proliferation of once-transformed and non-transformed plant cells of the same type into said first encrypted carbon source that supports growth and proliferation of said twice- and once-transformed cells, said second
30 gene being operatively linked to (ii) a second promoter DNA segment that controls expression of said second heterologous gene and (iii) a termination DNA segment;

(b) maintaining said heterotrophic culture
35 conditions for a time period sufficient for said twice- transformed plant cells to express said first

-83-

and second heterologous enzymes, grow and proliferate; and

(c) recovering said proliferating cells.

5 29. The process according to claim 28 wherein said second heterologous DNA selectable marker segment encodes the enzyme mannitol-1-oxo reductase, said second encrypted carbon source is mannitol, said first gene that encodes a heterologous gene encodes
10 phosphomannose isomerase and said first encrypted carbon source is mannose.

30. A process for selectively growing twice-transformed plants from a mixture of twice- and
15 once-transformed plant cells comprising the steps of:

(a) culturing a mixture of twice- and once-transformed plant cells under heterotrophic culture conditions in a culture medium that contains minimal
20 nutrients required for proliferation and growth by said once-transformed plant cells except for a source of carbon that supports growth and proliferation of said once-transformed cells and about 1.5 to 3 times the standard amount of
25 phosphorus, said source of carbon being replaced by a second encrypted carbon source that does not support growth and proliferation of said once-transformed plant cells; said twice-transformed cells containing first and a second heterologous DNA
30 segments that contains at least two expression cassettes, wherein at least one expression cassette is in the first heterologous DNA segment and at least one expression cassette is in the second heterologous DNA segment;
35 the expression cassette in the first hetero-

logous DNA segment containing a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression converts a first encrypted carbon source into a carbon source that supports growth and proliferation of said once- and twice-transformed plant cells under heterotrophic culture conditions but does not support growth and proliferation of non-transformed plant cells, said first heterologous gene being operatively linked to (ii) a first promoter DNA segment that controls expression of said first heterologous gene, and (iii) a termination DNA segment;

the expression cassette in the second heterologous DNA segment containing a second heterologous DNA selectable marker segment that includes (i) a second heterologous gene that encodes a second heterologous enzyme that on expression during heterotrophic culture of said twice-transformed cells converts said second encrypted carbon source that does not support growth and proliferation of once-transformed and non-transformed plant cells of the same type into said first encrypted carbon source that supports growth and proliferation of said twice- and once-transformed cells, said second heterologous gene being operatively linked to (ii) a second promoter DNA segment that controls expression of said second heterologous gene and (iii) a termination DNA segment;

(b) maintaining said heterotrophic culture conditions for a time period sufficient for said twice- transformed plant cells to express said first and second heterologous enzymes, grow and proliferate;

(c) recovering said proliferating cells; and
(d) regenerating plant meristematic tissues or plant embryos from said proliferating cells.

31. The process according to claim 30 wherein said second heterologous DNA selectable marker segment encodes the enzyme mannitol-1-oxo reductase, said second encrypted carbon source is mannitol, said first gene that encodes for the expression of phosphomannose isomerase and said first encrypted carbon source is mannose.

32. A kit for forming transformed plant cells comprising:

(a) a first package containing a DNA segment for transforming plant cells that contains an expression cassette operatively linked to a linker segment containing at least one restriction endonuclease site, said expression cassette containing a heterologous DNA selectable marker segment that includes (i) a first heterologous gene that encodes a heterologous enzyme that on expression during heterotrophic culture of transformed plant cells converts an encrypted carbon source that does not support growth and proliferation of non-transformed plant cells into a carbon source that supports growth and proliferation of said transformed cells, said first gene being operatively linked to (ii) a promoter DNA segment that controls expression of said first heterologous gene and (iii) a termination DNA segment; and

(b) a second package that contains minimal nutrients required for proliferation and growth of non-transformed plant cells during heterotrophic culture except for a source of carbon and about 1.5 to 3 times the standard amount of phosphorus, said source of carbon being replaced by an encrypted carbon source that does not support growth and proliferation of non-transformed plant cells but supports growth and proliferation of transformed

-86-

plant cells whose genome contains the DNA segment of said first package.

33. The kit according to claim 32 wherein said
5 heterologous gene is phosphomannose isomerase and said encrypted carbon source is mannose.

34. The kit according to claim 32 wherein said
10 first promoter DNA segment that controls expression of said heterologous gene is repressed by a product of the normal metabolism of said transgenic plants under autotrophic growth.

35. The kit according to claim 32 wherein said
15 linker segment contains a plurality of endonuclease restriction sites.


36. The kit according to claim 32 wherein said
20 DNA segment of (a) is located within the TDNA borders of the Ti plasmid of *Agrobacterium tumefaciens*.

37. The kit according to claim 32 wherein said
25 phosphorous is a water-soluble phosphate salt.

38. The kit according to claim 37 wherein the
phosphate salt is selected from the group consisting of: potassium phosphate, sodium phosphate, calcium phosphate, sodium hexametaphosphate, ammonium
30 phosphate, orthophosphoric acid, ferric phosphate, sodium glycerophosphate, ferric glycerophosphate and organic phosphate.


1/3

FIG. 1



2/3

FIG. 2



3/3

FIG. 3

NPTII
EXPRESSION